



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2500320 A1 2004/04/29

(21) 2 500 320

(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2003/10/09

(87) Date publication PCT/PCT Publication Date: 2004/04/29

(85) Entrée phase nationale/National Entry: 2005/03/24

(86) N° demande PCT/PCT Application No.: EP 2003/011203

(87) N° publication PCT/PCT Publication No.: 2004/035569

(30) Priorités/Priorities: 2002/10/10 (102 47 233.5) DE;
2002/12/02 (102 56 405.1) DE

(51) Cl.Int.⁷/Int.Cl.⁷ C07D 407/14, C07D 239/88,
C07D 239/72, C07D 451/04, C07D 491/04,
C07D 471/04, C07D 215/20, C07D 417/14,
C07D 413/14, C07D 417/12, C07D 413/12,
C07D 407/12, C07D 401/12, C07D 401/06


(71) Demandeur/Applicant:
MORPHOCHEM AKTIENGESELLSCHAFT FUER
KOMBINATORISCHE CHEMIE, DE

(72) Inventeurs/Inventors:
SURIVET, JEAN-PHILIPPE, FR;
ZUMBRUNN, CORNELIA, CH;
HUBSCHWERLEN, CHRISTIAN, FR;
PEREZ FRUTOS HOENER, ANNABELLE, CH

(74) Agent: MACRAE & CO.

(54) Titre : NOUVEAUX COMPOSES ANTIBACTERIENS

(54) Title: NOVEL COMPOUNDS WITH ANTIBACTERIAL ACTIVITY



(57) Abrégé/Abstract:

The invention relates to novel antibacterial compounds of formula (I).

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG(19) Weltorganisation für geistiges Eigentum
Internationales Büro(43) Internationales Veröffentlichungsdatum
29. April 2004 (29.04.2004)

PCT

(10) Internationale Veröffentlichungsnummer
WO 2004/035569 A3(51) Internationale Patentklassifikation⁷: C07D 407/14,
401/06, 413/14, 417/14, 407/12, 417/12, 239/72, 471/04,
401/12, 413/12, 491/04, 215/20, 239/88, 451/04(74) Anwälte: FORSTMAYER, Dietmar usw.; Boeters &
Lieck, Bereiteranger 15, 81541 München (DE).

(21) Internationales Aktenzeichen: PCT/EP2003/011203

(81) Bestimmungsstaaten (*national*): AE, AG, AL, AM, AT,
AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.(22) Internationales Anmeldedatum:
9. Oktober 2003 (09.10.2003)

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:
102 47 233.5 10. Oktober 2002 (10.10.2002) DE
102 56 405.1 2. Dezember 2002 (02.12.2002) DE(84) Bestimmungsstaaten (*regional*): ARIPO Patent (GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), europäisches Patent (AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL,
PT, RO, SE, SI, SK, TR), OAPI Patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).(71) Anmelder (*für alle Bestimmungsstaaten mit Ausnahme von
US*): MORPHOCHEM AKTIENGESELLSCHAFT
FÜR KOMBINATORISCHE CHEMIE [DE/DE];
Gmunder Strasse 37-37a, 81379 München (DE).

Veröffentlicht:


— mit internationalem Recherchenbericht

(72) Erfinder; und

(75) Erfinder/Anmelder (*nur für US*): SURIVET,
Jean-Philippe [FR/FR]; 19, rue Theo Bachmann, F-68300
Saint-Louis (FR). ZUMBRUNN, Cornelia [CH/CH];
Alemanngasse 57, CH-4058 Basel (CH). HUBSCHW-
ERLEN, Christian [FR/FR]; 15, rue de la Gendarmerie,
F-68480 Durmenach (FR). PÉREZ FRUTOS HÖNER,
Annabelle [CH/CH]; Mittlere Strasse 148, CH-4056 Basel
(CH).(88) Veröffentlichungsdatum des internationalen
Recherchenberichts: 2. September 2004Zur Erklärung der Zweibuchstaben-Codes und der anderen Ab-
kürzungen wird auf die Erklärungen ("Guidance Notes on Co-
des and Abbreviations") am Anfang jeder regulären Ausgabe der
PCT-Gazette verwiesen.

(54) Title: NOVEL COMPOUNDS WITH ANTIBACTERIAL ACTIVITY

(54) Bezeichnung: NEUE VERBINDUNGEN MIT ANTIBAKTERIELLER AKTIVITÄT



(I)

(57) Abstract: The invention relates to novel antibacterial compounds
of formula (I).(57) Zusammenfassung: Die vorliegende Anmeldung beschreibt neu-
artige antibakterielle Verbindungen der Formel (I).

WO 2004/035569 A3


Novel compounds having anti-bacterial activity

Resistance to the antibiotics used currently has increased appreciably in many countries of the world in recent years and in some cases has assumed alarming proportions. The main problem is that those pathogens exhibit not just a single resistance but, as a rule, multiple resistance. This is true especially of some gram-positive pathogen groups, such as staphylococci, pneumococci and enterococci (S. Ewig et al.; Antibiotika-Resistenz bei Erregern ambulant erworbener Atemwegsinfektionen (Antibiotic resistance in pathogens of outpatient-acquired respiratory tract infections); Chemother. J. 2002, 11, 12-26; F. Tenover; Development and spread of bacterial resistance to antimicrobial agents: an overview; Clin. Infect. Dis. 2001 Sep 15, 33 Suppl. 3, 108-115)

A long-feared development has recently occurred: In the USA, the first strain of *Staphylococcus aureus* has been described that is not only resistant to methicillin but also highly resistant to vancomycin (Centers for Disease Control and Prevention; *Staphylococcus aureus* resistant to vancomycin - United States, 2002; MMWR 2002, 51, 565-567). In addition to hygiene measures in hospitals, therefore, increased efforts are also required to find new antibiotics that as far as possible have a novel structure and a novel mechanism of action so as to be effective against those problem bacteria.

The present invention describes new kinds of compounds having anti-bacterial activity.

The present invention relates to compounds of the general formula (I):



wherein

5


A is an oxygen, sulphur or nitrogen atom or a C₁₋₄alkylene, C₂₋₄alkenylene, C₂₋₄alkynylene or C₁₋₄heteroalkylene group,

10 X₁, X₂, X₃, X₄ and X₅ are each independently of the others nitrogen atoms or groups of formula CR²,


R¹ is a hydrogen atom, a halogen atom, a hydroxy group, an alkyloxy group or a heteroalkyloxy group,


15 R² is a hydrogen atom, a halogen atom, or a hydroxy, alkyl, alkenyl, alkynyl or heteroalkyl group,

R³ is selected from the following groups:



20





the radicals R^4 , each independently of any other(s), are a hydroxy group, a C_{1-6} alkyl group or a C_{1-8} heteroalkyl group,

5

R^5 is an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkyl-cycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical,

10

n is 0, 1, 2 or 3 and

m is 0 or 2,

15 or a pharmacologically acceptable salt, solvate, hydrate or a pharmacologically acceptable formulation thereof.

The expression alkyl refers to a saturated, straight-chain or branched hydrocarbon group that contains from 1 to 20 carbon atoms, preferably from 1 to 12 carbon atoms, especially from 1 to 8 or from 1 to 6 or from 1 to 4 carbon atoms, for example a methyl, ethyl, propyl, isopropyl, isobutyl, *tert*-butyl, *n*-hexyl, 2,2-dimethyl-butyl or *n*-octyl group.

25

The expressions alkenyl and alkynyl refer to at least partially unsaturated, straight-chain or branched hydrocarbon groups that contain from 2 to 20 carbon atoms, preferably from 2 to 12 carbon atoms, especially from 2 to

6 carbon atoms, for example an ethenyl, allyl, acetylenyl, propargyl, isoprenyl or hex-2-enyl group. Preferably, alkenyl groups have one or two (especially one) double bond(s) and alkynyl groups have one or two (especially one) triple bond(s).

Furthermore, the terms alkyl, alkenyl and alkynyl refer to groups in which one or more hydrogen atoms have been replaced by a halogen atom (preferably F or Cl) such as, for example, a 2,2,2-trichloroethyl or a trifluoromethyl group.

The expression heteroalkyl refers to an alkyl, alkenyl or alkynyl group in which one or more (preferably 1, 2 or 3) carbon atoms have been replaced by an oxygen, nitrogen, phosphorus, boron, selenium, silicon or sulphur atom (preferably oxygen, sulphur or nitrogen). The expression heteroalkyl furthermore refers to a carboxylic acid or to a group derived from a carboxylic acid such as, for example, acyl, acylalkyl, alkoxycarbonyl, acyloxy, acyloxyalkyl, carboxyalkylamide or alkoxycarbonyloxy.

Examples of heteroalkyl groups are groups of formulae R^a-O-Y^a- , R^a-S-Y^a- , $R^a-N(R^b)-Y^a-$, R^a-CO-Y^a- , $R^a-O-CO-Y^a-$, $R^a-CO-O-Y^a-$, $R^a-CO-N(R^b)-Y^a-$, $R^a-N(R^b)-CO-Y^a-$, $R^a-O-CO-N(R^b)-Y^a-$, $R^a-N(R^b)-CO-O-Y^a-$, $R^a-N(R^b)-CO-N(R^c)-Y^a-$, $R^a-O-CO-O-Y^a-$, $R^a-N(R^b)-C(=NR^d)-N(R^c)-Y^a-$, R^a-CS-Y^a- , $R^a-O-CS-Y^a-$, $R^a-CS-O-Y^a-$, $R^a-CS-N(R^b)-Y^a-$, $R^a-N(R^b)-CS-Y^a-$, $R^a-O-CS-N(R^b)-Y^a-$, $R^a-N(R^b)-CS-O-Y^a-$, $R^a-N(R^b)-CS-N(R^c)-Y^a-$, $R^a-O-CS-O-Y^a-$, $R^a-S-CO-Y^a-$, $R^a-CO-S-Y^a-$, $R^a-S-CO-N(R^b)-Y^a-$, $R^a-N(R^b)-CO-S-Y^a-$, $R^a-S-CO-O-Y^a-$, $R^a-O-CO-S-Y^a-$, $R^a-S-CO-S-Y^a-$, $R^a-S-CS-Y^a-$, $R^a-CS-S-Y^a-$, $R^a-S-CS-N(R^b)-Y^a-$,

$R^a-N(R^b)-CS-S-Y^a-$, $R^a-S-CS-O-Y^a-$, $R^a-O-CS-S-Y^a-$, R^a being a hydrogen atom, a C_1-C_6 alkyl, a C_2-C_6 alkenyl or a C_2-C_6 alkynyl group; R^b being a hydrogen atom, a C_1-C_6 alkyl, a C_2-C_6 alkenyl or a C_2-C_6 alkynyl group; R^c being a hydrogen atom, a C_1-C_6 alkyl, a C_2-C_6 alkenyl or a C_2-C_6 alkynyl group; R^d being a hydrogen atom, a C_1-C_6 alkyl, a C_2-C_6 alkenyl or a C_2-C_6 alkynyl group and Y^a being a direct bond, a C_1-C_6 alkylene, a C_2-C_6 alkenylene or a C_2-C_6 alkynylene group, each heteroalkyl group containing at least one carbon atom and it being possible for one or more hydrogen atoms to have been replaced by fluorine or chlorine atoms. Specific examples of heteroalkyl groups are methoxy, trifluoromethoxy, ethoxy, n-propyloxy, isopropyloxy, tert-butyloxy, methoxymethyl, ethoxymethyl, methoxyethyl, methylamino, ethylamino, dimethylamino, diethylamino, isopropylethylamino, methylaminomethyl, ethylaminomethyl, diisopropylaminoethyl, enol ether, dimethylaminomethyl, dimethylaminoethyl, acetyl, propionyl, butyryloxy, acetyloxy, methoxycarbonyl, ethoxycarbonyl, N-ethyl-N-methylcarbamoyl and N-methylcarbamoyl. Further examples of heteroalkyl groups are nitrile, isonitrile, cyanate, thiocyanate, isocyanate, isothiocyanate and alkyl nitrile groups.

The expression cycloalkyl refers to a saturated or partially unsaturated (e.g. cycloalkenyl) cyclic group that contains one or more rings (preferably 1 or 2) containing from 3 to 14 ring carbon atoms, preferably from 3 to 10 (especially 3, 4, 5, 6 or 7) ring carbon atoms. The expression cycloalkyl refers furthermore to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O,

SH, =S, NH₂, =NH or NO₂ groups, thus, for example, cyclic ketones such as, for example, cyclohexanone, 2-cyclohexenone or cyclopentanone. Further specific examples of cycloalkyl groups are a cyclopropyl, cyclobutyl, cyclopentyl, spiro[4,5]decanyl, norbornyl, cyclohexyl, cyclopentenyl, cyclohexadienyl, decaliny, cubanyl, bicyclo[4.3.0]nonyl, tetralin, cyclopentylcyclohexyl, fluorocyclohexyl or cyclohex-2-enyl group.

10 The expression heterocycloalkyl refers to a cycloalkyl group as defined above in which one or more (preferably 1, 2 or 3) ring carbon atoms have been replaced by an oxygen, nitrogen, silicon, selenium, phosphorus or sulphur atom (preferably oxygen, sulphur or nitrogen). A heterocycloalkyl group has preferably 1 or 2 ring(s) containing from 3 to 10 (especially 3, 4, 5, 6 or 7) ring atoms. The expression heterocycloalkyl refers furthermore to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH or NO₂ groups. Examples are a piperidyl, morpholinyl, urotropinyl, pyrrolidinyl, tetrahydrothiophenyl, tetrahydropyranyl, tetrahydrofuryl, oxacyclopentyl, azacyclopentyl or 2-pyrazolinyl group and also lactams, lactones, cyclic imides and cyclic anhydrides.

25

The expression alkylcycloalkyl refers to groups containing both cycloalkyl and also alkyl, alkenyl or alkynyl groups in accordance with the above definitions, for example alkylcycloalkyl, alkylcycloalkenyl, alkenylcycloalkyl and alkynylcycloalkyl groups. An alkylcycloalkyl group preferably contains a cycloalkyl group that contains one or two rings systems having from 3 to 10 (especially 3, 4,

5, 6 or 7) ring carbon atoms, and one or two alkyl, alkenyl or alkynyl groups having 1 or 2 to 6 carbon atoms.

The expression heteroalkylcycloalkyl refers to alkylcyclo-
5 alkyl groups as defined above in which one or more
(preferably 1, 2 or 3) carbon atoms have been replaced by
an oxygen, nitrogen, silicon, selenium, phosphorus or
sulphur atom (preferably oxygen, sulphur or nitrogen). A
heteroalkylcycloalkyl group preferably contains 1 or 2
10 ring systems having from 3 to 10 (especially 3, 4, 5, 6 or
7) ring atoms, and one or two alkyl, alkenyl, alkynyl or
heteroalkyl groups having 1 or 2 to 6 carbon atoms.
Examples of such groups are alkylheterocycloalkyl, alkyl-
heterocycloalkenyl, alkenylheterocycloalkyl, alkynyl-
15 heterocycloalkyl, heteroalkylcycloalkyl, heteroalkyl-
heterocycloalkyl and heteroalkylheterocycloalkenyl, the
cyclic groups being saturated or mono-, di- or tri-
unsaturated.

20 The expression aryl or Ar refers to an aromatic group that
has one or more rings containing from 6 to 14 ring carbon
atoms, preferably from 6 to 10 (especially 6) carbon
atoms. The expression aryl (or Ar) refers furthermore to
groups in which one or more hydrogen atoms have been
25 replaced by fluorine, chlorine, bromine or iodine atoms or
by OH, SH, NH₂ or NO₂ groups. Examples are a phenyl,
naphthyl, biphenyl, 2-fluorophenyl, aniliny, 3-nitro-
phenyl or 4-hydroxyphenyl group.

30 The expression heteroaryl refers to an aromatic group that
has one or more rings and is formed by a ring system that
contains from 5 to 14 ring atoms, preferably from 5 to 10

(especially 5 or 6) ring atoms, and contains one or more (preferably 1, 2, 3 or 4) oxygen, nitrogen, phosphorus or sulphur ring atoms (preferably O, S or N). The expression heteroaryl refers furthermore to groups in which one or
5 more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, SH, NH₂ or NO₂ groups. Examples are 4-pyridyl, 2-imidazolyl, 3-phenyl-pyrrolyl, thiazolyl, oxazolyl, triazolyl, tetrazolyl, isoxazolyl, indazolyl, indolyl, benzimidazolyl, pyrid-
10 azinyl, quinolinyl, purinyl, carbazolyl, acridinyl, pyrimidyl, 2,3'-bifuryl, 3-pyrazolyl and isoquinolinyl groups.

The expression aralkyl refers to groups containing both
15 aryl and also alkyl, alkenyl, alkynyl and/or cycloalkyl groups in accordance with the above definitions, such as, for example, arylalkyl, arylalkenyl, arylalkynyl, aryl-cycloalkyl, arylcycloalkenyl, alkylarylcycloalkyl and alkylarylcycloalkenyl groups. Specific examples of
20 aralkyls are toluene, xylene, mesitylene, styrene, benzyl chloride, o-fluorotoluene, 1H-indene, tetralin, dihydronaphthalene, indanone, phenylcyclopentyl, cumene, cyclohexylphenyl, fluorene and indan. An aralkyl group preferably contains one or two aromatic ring systems (1 or
25 2 rings) containing from 6 to 10 carbon atoms and one or two alkyl, alkenyl and/or alkynyl groups containing 1 or 2 to 6 carbon atoms and/or a cycloalkyl group containing 5 or 6 ring carbon atoms.

30 The expression heteroaralkyl refers to an aralkyl group as defined above in which one or more (preferably 1, 2, 3 or 4) carbon atoms have been replaced by an oxygen, nitrogen,

silicon, selenium, phosphorus, boron or sulphur atom (preferably oxygen, sulphur or nitrogen), that is to say to groups containing both aryl or heteroaryl and also alkyl, alkenyl, alkynyl and/or heteroalkyl and/or cycloalkyl and/or heterocycloalkyl groups in accordance with the above definitions. A heteroaralkyl group preferably contains one or two aromatic ring systems (1 or 2 rings) containing 5 or 6 to 10 carbon atoms and one or two alkyl, alkenyl and/or alkynyl groups containing 1 or 2 to 6 carbon atoms and/or a cycloalkyl group containing 5 or 6 ring carbon atoms, 1, 2, 3 or 4 or those carbon atoms having been replaced by oxygen, sulphur or nitrogen atoms.

Examples are arylheteroalkyl, arylheterocycloalkyl, arylheterocycloalkenyl, arylalkylheterocycloalkyl, arylalkenylheterocycloalkyl, arylalkynylheterocycloalkyl, arylalkylheterocycloalkenyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, heteroarylheteroalkyl, heteroarylcycloalkyl, heteroarylcycloalkenyl, heteroarylheterocycloalkyl, heteroarylheterocycloalkenyl, heteroarylalkylcycloalkyl, heteroarylalkylheterocycloalkenyl, heteroarylheteroalkylcycloalkyl, heteroarylheteroalkylcycloalkenyl and heteroarylheteroalkylheterocycloalkyl groups, the cyclic groups being saturated or mono-, di- or tri-unsaturated. Specific examples are a tetrahydroisoquinolyl-, benzoyl-, 2- or 3-ethylindolyl-, 4-methylpyridino-, 2-, 3- or 4-methoxyphenyl-, 4-ethoxyphenyl-, 2-, 3- or 4-carboxyphenyl-alkyl group.

The expressions cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aryl, heteroaryl, aralkyl and heteroaralkyl refer to groups in which one or


more hydrogen atoms of such groups have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH or NO₂ groups.

- 5 The expression "optionally substituted" refers to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH or NO₂ groups. The expression refers furthermore to groups that are substituted by
- 10 unsubstituted C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆heteroalkyl, C₃-C₁₀cycloalkyl, C₂-C₉heterocycloalkyl, C₆-C₁₀aryl, C₁-C₉heteroaryl, C₇-C₁₂aralkyl or C₂-C₁₁hetero-aralkyl groups.
- 15 Owing to their substitution, compounds of formula (I) may contain one or more centres of chirality. The present invention therefore includes both all pure enantiomers and all pure diastereoisomers and also mixtures thereof in any mixing ratio. The present invention moreover also
- 20 includes all *cis/trans*-isomers of the compounds of the general formula (I) and also mixtures thereof. The present invention moreover includes all tautomeric forms of the compounds of formula (I).
- 25 Preferred are compounds of formula (I) wherein A is an oxygen atom or a group of formula CH₂ or CH(OH).

Also preferred are compounds of formula (I) wherein one of the groups X₁, X₂, X₃, X₄ and X₅ (especially X₁ or X₂) is a

30 nitrogen atom and the others are CH groups, or all of the groups X₁, X₂, X₃, X₄ and X₅ are CH groups.

Furthermore, R^3 is preferably one of the following groups:



R^1 is especially preferably a halogen atom, a C_{1-6} alkyloxy group (for example methoxy or ethoxy), a methyl group or an ethyl group; especially a methoxy group.

Also preferably, R^4 is a C_{1-6} heteroalkyl group, as defined hereinabove, having one or two oxygen atoms as individual hetero atoms.

R^4 is especially preferably a group of formula $-COOH$, $-CH_2COOH$, $-CH_2CH_2COOH$, $-CH_2COOCH_3$, $-CH_2CH_3$, $-CH_2OH$, $-CH_2CH_2OH$, $-OH$, $-OCH_3$, $-CH_2OCONH_2$, $-CH_2CH_2COOCH_3$, $-COOCH_3$, $-CH_3$ or $-(CH_2)_3OH$.

Also preferably, n is 0 or 1.

Also preferably, R^5 is an aralkyl or a hetero-aralkyl group.

Again preferably, R^5 is a group of formula $-Y-Cy$, Y being a C_1-C_6 alkylene, C_2-C_6 alkenylene or C_1-C_6 heteroalkylene group, wherein optionally a hydrogen atom may have been replaced by a hydroxy group or two hydrogen atoms may have been replaced by an $=O$ group, and Cy being an optionally substituted phenyl, naphthyl or heteroaryl group containing 1 or 2 rings and from 5 to 10 ring atoms, or an optionally substituted arylheterocycloalkyl or heteroaryl-

heterocycloalkyl group containing two rings and 9 or 10 ring atoms.

The therapeutic use of compounds of formula (I), their
5 pharmacologically acceptable salts or solvates and hydrates and also formulations and pharmaceutical compositions also lie within the scope of the present invention.

10 The pharmaceutically compositions according to the present invention comprise at least one compound of formula (I) as active ingredient and, optionally, carrier substances and/or adjuvants.

15 Examples of pharmacologically acceptable salts of the compounds of formula (I) are salts of physiologically acceptable mineral acids, such as hydrochloric acid, sulphuric acid and phosphoric acid, and salts of organic acids, such as methanesulphonic acid, p-toluenesulphonic
20 acid, lactic acid, acetic acid, trifluoroacetic acid, citric acid, succinic acid, fumaric acid, oxalic acid, maleic acid and salicylic acid. Further examples of pharmacologically acceptable salts of the compounds of formula (I) are alkali metal and alkaline earth metal
25 salts such as, for example, sodium, potassium, lithium, calcium and magnesium salts, ammonium salts and salts of organic bases such as, for example, methylamine, dimethylamine, triethylamine, piperidine, ethylenediamine, lysine, choline hydroxide, meglumine, morpholine and
30 arginine salts. Compounds of formula (I) can be solvated, especially hydrated. The hydration may take place, for example, during the preparation process or as a

consequence of the hygroscopic nature of the initially anhydrous compounds of formula (I). When the compounds of formula (I) comprise asymmetric carbon atoms, they may be either in the form of achiral compounds, diastereoisomeric mixtures, mixtures of enantiomers or in the form of optically pure compounds.

The pro-drugs to which the present invention also relates consist of a compound of formula (I) and at least one pharmacologically acceptable protecting group that is removed under physiological conditions, for example an alkoxy, aralkyloxy, acyl or acyloxy group, such as, for example, an ethoxy, daloxate, benzyloxy, acetyl or acetyloxy group or a group of formula $\text{COOCH}_2\text{OCOC}(\text{CH}_3)_3$ or $\text{COOCH}_2\text{OCOO-cyclohexyl}$.

The present invention relates also to the use of those active ingredients in the preparation of medicaments. In general, compounds of formula (I) are administered either individually, or in combination with any other desired therapeutic agent, using the known and acceptable methods. Such therapeutically useful agents may be administered, for example, by one of the following routes: orally, for example in the form of dragées, coated tablets, pills, semi-solid substances, soft or hard capsules, solutions, emulsions or suspensions; parenterally, for example in the form of an injectable solution; rectally in the form of suppositories; by inhalation, for example in the form of a powder formulation or spray, transdermally or intranasally. For the preparation of such tablets, pills, semi-solid substances, coated tablets, dragées and hard gelatin capsules, the therapeutically usable product can

be mixed with pharmacologically inert, inorganic or organic pharmaceutical carrier substances, for example with lactose, sucrose, glucose, gelatin, malt, silica gel, starch or derivatives thereof, talcum, stearic acid or salts thereof, skimmed milk powder and the like. For the preparation of soft capsules, pharmaceutical carrier substances such as, for example, vegetable oils, petroleum, animal or synthetic oils, wax, fat and polyols can be used. For the preparation of liquid solutions and syrups, pharmaceutical carrier substances such as, for example, water, alcohols, aqueous saline solution, aqueous dextrose, polyols, glycerol, vegetable oils, petroleum and animal or synthetic oils can be used. For suppositories, pharmaceutical carrier substances such as, for example, vegetable oils, petroleum, animal or synthetic oils, wax, fat and polyols can be used. For aerosol formulations, compressed gases that are suitable for the purpose can be used, such as, for example, oxygen, nitrogen and carbon dioxide. The pharmaceutically acceptable agents may also comprise additives for preserving and stabilising, emulsifiers, sweeteners, flavourings, salts for altering the osmotic pressure, buffers, encapsulation additives and anti-oxidants.

Combinations with other therapeutic agents may comprise other antimicrobial and anti-fungal active ingredients.

For the prevention and/or treatment of bacterial infections, the dose of the biologically active compound according to the invention can vary within wide limits and can be adjusted to individual requirements. Generally, a dose of from 10 mg to 4000 mg per day is suitable, a


preferred dose being from 50 to 3000 mg per day. In suitable cases, the dose may also be below or above the stated values. The daily dose can be administered as a single dose or in a plurality of doses. A typical individual dose contains approximately 50 mg, 100 mg, 250 mg, 500 mg, 1 g or 2 g of the active ingredient.

EXAMPLES

10

Example 1: 2-((3RS)-{[(2,3-Dihydro-benzo[1,4]dioxin-6-yl-methyl)amino]methyl}piperidin-1-yl)-(1RS)-(6-methoxy-quinolin-4-yl)ethanol.

15 1.a) Benzyl (3RS)-azidomethylpiperidine-1-carbamate:



At 0°C, triethylamine (5.6 ml, 40.1 mmol) and then methanesulphonyl chloride (2 ml, 25.7 mmol) were added to a solution of benzyl 3-hydroxymethylpiperidine-1-carbamate (5 g, 20.05 mmol) in dichloromethane (100 ml). After the solution had been stirred for 20 min., the reaction mixture was cooled to -60°C and a solution of benzyl 4-oxopiperidine-1-carbamate (2.33 g, 10 mmol) in diethyl ether (10 ml) was added. The reaction mixture was heated to room temperature in the course of 30 minutes and water (40 ml) was added. The two phases were separated and the aqueous phase was extracted twice using 50 ml of ethyl


acetate each time. The combined organic phases were washed with 20 ml of saturated sodium chloride solution, dried over MgSO_4 , filtered and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (EtOAc). The resulting oil was dissolved in DMF (45 ml), and sodium azide (2.6 g, 40 mmol) was added. The reaction mixture was further stirred for 3 hours at 80°C and then diluted with water (100 ml) and ethyl acetate (200 ml). The two phases were separated and the aqueous phase was extracted twice using 50 ml of ethyl acetate each time. The combined organic phases were washed with 20 ml of saturated sodium chloride solution, dried over MgSO_4 , filtered and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (hexane/EtOAc 1/1).

Yield: 6.1 g (18.6 mmol)

^1H -NMR (CDCl_3 , 300 MHz): 1.28 (m, 1H); 1.51 (m, 1H); 1.60-1.87 (m, 3H); 2.74 (br s, 1H); 2.91 (m, 1H); 3.23 (br d, $J = 4.5$ Hz, 2H); 3.98 (td, $J = 4.1, 13.2$ Hz, 1H); 4.06 (br s, 1H); 5.15 (s, 2H); 7.28-7.38 (m, 5H).

Benzyl 3-hydroxymethyl-piperidine-1-carbamate has already been described in *Arch. Pharm.* (Weinheim, Germany) 1990 p. 9-12.

1.b) Benzyl (3RS)-aminomethyl-piperidine-1-carbamate:




Triphenylphosphine (8 g, 30 mmol) was added to a solution of benzyl 3-azidomethylpiperidine-1-carbamate (6.1 g,

18.6 mmol) in THF (37 ml) and water (5 ml). After the solution had been stirred for 3 hours at 60°C, the reaction mixture was concentrated and the residue was taken up in 3N HCl (200 ml) and ether (200 ml). The two
 5 phases were separated and the aqueous phase was extracted twice using 100 ml of ethyl acetate each time. Solid sodium hydroxide (16 g, 640 mmol) was cautiously added until an oil separated out. The mixture was diluted with ethyl acetate and the organic phases was dried over MgSO₄
 10 and filtered, and the filtrate was concentrated to dryness by rotary evaporation.

Yield: 2.81 g, 11.3 mmol

MS (EI) m/z 249 [M+H]⁺

15 1.c) Benzyl (3RS)-{[(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-amino]-methyl}-piperidine-1-carbamate:




20 1,4-Benzodioxane-6-carbaldehyde (0.984 g, 6 mmol) and sodium triacetoxyborohydride (1.7 g, 8 mmol) were added to a solution of benzyl 3-aminomethylpiperidine-1-carbamate (1.5 g, 6 mmol) in dichloroethane (37 ml) and THF (4 ml). After the solution had been stirred at room temperature
 25 for 3 hours, saturated NaHCO₃ solution (20 ml) was added. The two phases were separated and the aqueous phase was extracted twice using 50 ml of dichloromethane each time. The combined organic phases were washed with 20 ml of saturated sodium chloride solution, dried over MgSO₄,
 30 filtered and concentrated to dryness by rotary

evaporation. The residue was purified by column chromatography on silica gel (EtOAc and then MeOH/EtOAc 1/9).

Yield: 1.7 g, 4.28 mmol (oil)

5 MS (EI) m/z 397 $[M+H]^+$

1.d (2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-piperidin-(3RS)-yl-methylamine:



10


20% $Pd(OH)_2$ on carbon (0.23 g) was added to a solution of benzyl 3-[[[(2,3-dihydrobenzo[1,4]dioxin-6-yl-methyl)-amino]methyl]piperidine-1-carbamate (0.98 g, 2.47 mmol) in EtOH (10 ml) and EtOAc (10 ml) and hydrogenation was
15 carried out under a hydrogen atmosphere (1 bar). The reaction mixture was filtered and the filtrate was concentrated.

Yield: 0.64 g, 2.43 mmol

MS (EI) m/z 308 $[M+H]^+$

20

1.e) (RS)-6-Methoxy-4-oxiranylquinoline:



Trimethylsulphonium iodide (0.954 g, 4.67 mmol) and
25 potassium hydroxide powder (1.8 g, 32 mmol) were added to a solution of 6-methoxyquinoline-4-carbaldehyde (0.85 g, 4.54 mmol) in acetonitrile (13.5 ml) and water (6 drops).


The reaction mixture was heated at 60°C for one hour. The mixture was then cooled to room temperature and benzene (40 ml) was added. The precipitate was filtered off and the filtrate was concentrated to dryness by rotary evaporation. The residue was diluted with water (100 ml) and ethyl acetate (100 ml), the phases were separated, and the aqueous phase was extracted twice using 50 ml of ethyl acetate each time. The combined organic phases were dried over MgSO₄, filtered and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (EtOAc).

Yield: 0.904 g, 4.5 mmol

MS (EI) m/z 202 [M+H]⁺

6-Methoxyquinoline-4-carbaldehyde was prepared according to *Eur. J. Med. Chem. Chim. Ther.* 2000 (35) p-707-714.

1.f) 2-((3RS)-{[(2,3-Dihydrobenzo[1,4]dioxin-6-ylmethyl)-amino]-methyl}-piperidin-1-yl)-(1RS)-(6-methoxyquinolin-4-yl)ethanol:



A solution of (RS)-6-methoxy-4-oxiranylquinoline (0.161 g, 0.8 mmol) and (2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-piperidin-(3RS)-ylmethylamine (0.210 g, 0.8 mmol) in ethanol (4 ml) was heated at 80°C for 16 hours. The reaction mixture was cooled to room temperature and concentrated to dryness by rotary evaporation. The residue

was purified by column chromatography on silica gel (EtOAc and then EtOAc/MeOH 4:1).


Yield: 0.240 g. 0.51 mmol (oil)

MS (EI) m/z 464 [M+H]⁺

5

Example 2: (1RS)-(6-Methoxy-quinolin-4-yl)-2-((3RS)-
{[(naphth-2-ylmethyl)-amino]-methyl}-piperidin-1-yl)-
ethanol

10 2.a) Benzyl (3RS)-(tert-butoxycarbonylamino-methyl)-
 piperidine-1-carbamate




Triethylamine (5.6 ml) and di-tert-butyl dicarbonate (4.9 g, 22.4 mmol) were added to a solution of benzyl 3-
 15 aminomethylpiperidine-1-carbamate (5 g, 20.1 mmol) in
 dichloromethane (100 ml). After the solution had been
 stirred at room temperature for 4 hours, the solution was
 concentrated to dryness by rotary evaporation. The residue
 was purified by column chromatography on silica gel
 20 (EtOAc/hexane 1/4).

Yield: 5.0 g, 14.3 mmol (oil)

MS (EI) m/z 349 [M+H]⁺

2.b) tert-Butyl (RS)-piperidin-3-ylmethylcarbamate




25

20% Pd(OH)₂ on carbon (1 g) was added to a solution of benzyl 3-(tert-butoxycarbonylamino-methyl)-piperidine-1-carbamate (5 g, 14.3 mmol) in EtOH (50 ml) and EtOAc (50 ml), and hydrogenation was carried out under a hydrogen atmosphere (1 bar). The reaction mixture was filtered and the filtrate was concentrated.

Yield: 3.0 g, 14 mmol (oil)

MS (EI) m/z 215 [M+H]⁺

- 10 2.c) *tert*-Butyl (1-[(2*RS*)-hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-piperidin-(3*RS*)-ylmethyl)carbamate




- A solution of 6-methoxy-4-oxiranylquinoline (2.68 g, 13.3 mmol) and *tert*-butyl piperidin-3-ylmethylcarbamate (2.85 g, 13.3 mmol) in ethanol (35 ml) was heated at 80°C for 12 hours. The reaction mixture was cooled to room temperature and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 9:1).

- 20 Yield: 2.49 g. 6 mmol (foam)

MS (EI) m/z 416 [M+H]⁺

- 2.d) 2-[(3*RS*)-Aminomethyl-piperidin-1-yl]-[(1*RS*)-(6-methoxy-quinolin-4-yl)-ethanol




A solution of *tert*-butyl {1-[2-hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-piperidin-3-ylmethyl}carbamate (2.49 g, 6 mmol) in TFA (10 ml) was stirred at room temperature for 20 minutes. The reaction mixture was concentrated and taken up in aqueous 2N NaOH. The aqueous phase was extracted with dichloromethane/MeOH (9/1), and the combined organic phases were dried over sodium sulphate, filtered and concentrated to dryness by rotary evaporation.

Yield: 1.7 g. 5.4 mmol (foam)

MS (EI) m/z 316 $[M+H]^+$

2.e) (1*RS*)-(6-Methoxy-quinolin-4-yl)-2-((3*RS*)-{[(naphth-2-ylmethyl)-amino]-methyl}-piperidin-1-yl)-ethanol




Molecular sieve, type 3A, (1 g) and 2-naphthaldehyde (0.078 g, 0.5 mmol) were added to a solution of 2-(3-aminomethyl-piperidin-1-yl)-1-(6-methoxy-quinolin-4-yl)-ethanol (0.158 g, 0.5 mmol) in MeOH (1.5 ml) and dichloromethane (3.5 ml). The reaction mixture was stirred at room temperature for 16 hours and sodium borohydride (0.05 g, 1.35 mmol) was added. The reaction mixture was stirred at room temperature for 2 hours and then filtered over Hydromatrix (wetted with a NaHCO_3 solution, 2 ml) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 9/1 1% NH_4OH).

Yield: 0.158 g. (0.34 mmol) (foam)


MS (EI) m/z 456 [M+H]⁺

5 Example 3: (1RS)-(6-Methoxy-quinolin-4-yl)-2-((3RS)-[(3-phenyl-allylamino)-methyl]-piperidin-1-yl)-ethanol



In an analogous manner, starting from Example 2.e, (1RS)-(6-methoxy-quinolin-4-yl)-(2RS)-{3-[(3-phenyl-allylamino)-methyl]-piperidin-1-yl}-ethanol was produced in a yield of 74% (MS (EI) m/z 432 [M+H]⁺).

Example 4: 2-((3RS)-[(3-Furan-2-yl-allylamino)-methyl]-piperidin-1-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol




15

In an analogous manner, starting from Example 2.e, 2-{3-[(3-furan-2-yl-allylamino)-methyl]-piperidin-1-yl}-1-(6-methoxy-quinolin-4-yl)-ethanol was produced in a yield of 56% (MS (EI) m/z 422 [M+H]⁺).

20

Example 5: (3S)-6-[[[1-[2-Hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-piperidin-(3R)-ylmethyl]-amino]-methyl]-4H-benzo[1,4]oxazin-3-one

5 5 a) *tert*-Butyl (3S)-aminomethyl-piperidine-1-carbamate




Triphenylphosphine (3.93 g, 15 mmol) was added to a solution of *tert*-butyl (3R)-azidomethyl-piperidine-1-carbamate (2.16 g, 8.9 mmol) in THF (60 ml) and water
 10 (5 ml). After the solution had been stirred at 60°C for 3 hours, the reaction mixture was concentrated and the residue was taken up in 3N HCl (200 ml) and ether (200 ml). The two phases were separated and the aqueous phase was extracted twice using 100 ml of ethyl acetate
 15 each time. Solid sodium hydroxide (6 g, 150 mmol) was cautiously added until an oil separated out. The mixture was diluted with ethyl acetate, the organic phase was dried over MgSO₄ and filtered and the filtrate was concentrated to dryness by rotary evaporation.

20 Yield: 1.90 g, 8.8 mmol

MS (EI) *m/z* 215 [M+H]⁺

tert-Butyl 3R-azidomethyl-piperidine-1-carbamate was prepared according to *J. Med. Chem.* **1994** (37) p. 3889-3901

5.b) *tert*-Butyl 3S-(benzyloxycarbonylamino-methyl)-piperidine-1-carbamate




Sodium bicarbonate (1.3 g, 15.4 mmol) and chloroformic acid benzyl ester (1.3 ml, 9 mmol) were added to a solution of *tert*-butyl 3S-aminomethyl-piperidine-1-carbamate (1.90 g, 8.8 mmol) in acetone/water (1/1, 100 ml). The reaction mixture was stirred for one hour and then concentrated to dryness by rotary evaporation. The residue was diluted with EtOAc (100 ml). The two phases were separated and the aqueous phase was extracted twice using 50 ml of EtOAc each time. The combined organic phases were washed with 20 ml of saturated sodium chloride solution, dried over MgSO₄, filtered and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (EtOAc/hexane 1/2).

Yield: 2.96 g , 8.5 mmol (oil)

MS (EI) m/z 349 [M+H]⁺

5.c) Benzyl piperidin-(3R)-ylmethylcarbamate




A solution of *tert*-butyl 3S-(benzyloxycarbonylamino-methyl)-piperidine-1-carbamate (2.96 g, 6 mmol) in TFA (10 ml) was stirred at room temperature for 20 minutes. The reaction mixture was concentrated and taken up in aqueous 2N NaOH. The aqueous phase was extracted with

dichloromethane/MeOH (9/1), and the combined organic phases were dried over sodium sulphate, filtered and concentrated to dryness by rotary evaporation.

Yield: 1.96 g. 7.9 mmol (foam)

5 MS (EI) m/z 249 [M+H]⁺

5.d) Benzyl (1-[(2RS)-hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-piperidin-(3R)-ylmethyl)carbamate




10 Potassium carbonate (1.12 g, 8.10 mmol) and lithium perchlorate (0.627 g, 5.89 mmol) were added to a solution of 6-methoxy-4-oxiranylquinoline (1.127 g, 5.60 mmol) and benzyl piperidin-3-ylmethylcarbamate (1.46 g, 5.88 mmol) in DMF (17 ml). The reaction mixture was heated at 60°C
15 for 16 hours and then cooled to room temperature and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 9:1 1% ammonium hydroxide).

Yield: 2.51 g. 5.58 mmol (oil)

20 MS (EI) m/z 450 [M+H]⁺

5.e) (3S)-2-(3-Aminomethyl-piperidin-1-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol




20% Pd(OH)₂ on carbon (0.7 g) was added to a solution of benzyl (1-[2-hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-piperidin-3-ylmethyl)carbamate (2.51 g, 5.58 mmol) in THF (35 ml), MeOH (7 ml) was added and hydrogenation was carried out under a hydrogen atmosphere (1 bar). The reaction mixture was filtered and the filtrate was concentrated.

Yield: 1.76 g. 5.58 mmol (oil)

MS (EI) m/z 316 [M+H]⁺

10

5.f) (3S)-6-[(1-[(2RS)-Hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-piperidin-3-ylmethyl)-amino)-methyl]-4H-benzo[1,4]oxazin-3-one



15 Molecular sieve, type 3A, (1.082 g) and 3-oxo-3,4-dihydro-2H-benzo[1,4]oxazine-6-carbaldehyde (0.095 g, 0.54 mmol) were added to a solution of (3S)-2-(3-aminomethyl-piperidin-1-yl)-1-(6-methoxy-quinolin-4-yl)-ethanol (0.171 g, 0.54 mmol) in MeOH (1.6 ml) and dichloromethane (3.8 ml). The reaction mixture was stirred at room temperature for 16 hours and sodium borohydride (0.054 g, 1.43 mmol) was added. The reaction mixture was stirred at room temperature for 2 hours and then filtered over Hydromatrix (wetted with an NaHCO₃ solution, 2 ml) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 9/1 1% NH₄OH).

20


25

Yield: 0.112 g. (0.24 mmol) (foam)

MS (EI) m/z 477 [M+H]⁺

3-Oxo-3,4-dihydro-2H-benzo[1,4]oxazine-6-carbaldehyde was
5 prepared according to WO 02/34754.


Example 6: (3S)-2-(3-(((Benzo[1,2,5]thiadiazol-5-ylmethyl)-amino)-methyl)-piperidin-1-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol



10

Starting from (3S)-2-(3-aminomethyl-piperidin-1-yl)-1-(6-methoxy-quinolin-4-yl)-ethanol (Example 5.e), (3S)-2-(3-
15 {{{benzo[1,2,5]thiadiazol-5-ylmethyl)-amino}-methyl}-piperidin-1-yl)-1-(6-methoxy-quinolin-4-yl)-ethanol was
produced in an analogous manner in a yield of 49% (MS (EI) m/z 464 [M+H]⁺).

Example 7: 2-((3S)-(((2,3-Dihydro-benzo[1,4]dioxepin-6-ylmethyl)-amino)-methyl)-piperidin-1-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol



20


Starting from (3S)-2-(3-aminomethyl-piperidin-1-yl)-1-(6-methoxy-quinolin-4-yl)-ethanol (Example 5.e), 2-((3S)-{[(2,3-dihydro-benzo[1,4]dioxepin-6-ylmethyl)-amino]-methyl}-piperidin-1-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol was produced in an analogous manner in a yield of 62% (MS (EI) m/z 478 [M+H]⁺).

3,4-Dihydro-2H-benzo[b][1,4]dioxepine-7-carbaldehyde was prepared according to *Chem. Abstr* **1958**, 3816.

10

Example 8: 2-((3S)-{[(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-methyl}-piperidin-1-yl)-(1S)-(6-methoxy-quinolin-4-yl)-ethanol

15 8.a) *tert*-Butyl (3S)-{[(2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-methyl}-piperidine-1-carbamate




Molecular sieve, type 3A, (9.6 g) and 2,3-dihydro-benzo[1,4]dioxin-6-carbaldehyde (0.078 g, 0.5 mmol) were added to a solution of *tert*-butyl 3S-aminomethyl-piperidine-1-carbamate (Example 5.a) (0.158 g, 0.5 mmol) in MeOH (8 ml) and dichloromethane (27 ml). The reaction mixture was stirred at room temperature for 16 hours and sodium borohydride (0.05 g, 1.35 mmol) was added. The reaction mixture was stirred at room temperature for 2 hours and then filtered over Hydromatrix (wetted with an

NaHCO₃ solution, 20 ml) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 19/1).

5 Yield: 2.2 g. 6.0 mmol (foam)

MS (EI) m/z 363.4 [M+H]⁺

8.b) (R)-(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-piperidin-3-ylmethyl-amine




10

A solution of *tert*-butyl 3S-([(2,3-dihydro-benzo[1,4]-dioxin-6-ylmethyl)-amino]-methyl)-piperidine-1-carbamate (2.2 g, 6 mmol) in TFA (10 ml) was stirred at room temperature for 20 minutes. The reaction mixture was concentrated and taken up in aqueous 2N NaOH. The aqueous phase
15 was extracted with dichloromethane/MeOH (9/1), and the combined organic phases were dried over sodium sulphate, filtered and concentrated to dryness by rotary evaporation.

20 Yield: 1.18 g, 4.53 mmol (foam)

MS (EI) m/z 263.4 [M+H]⁺

8.c) 2-((3S)-{[(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-methyl}-piperidin-1-yl)-(1S)-(6-methoxy-quinolin-4-yl)-ethanol




5 The title compound was produced in a manner analogous to Example 1.f in a yield of 36% (MS (EI) m/z 464.5 [M+H]⁺).

(R)-6-Methoxy-4-oxiranylquinoline was prepared analogously to WO 02/50040.

10

Example 9: 2-((3S)-{[(2,3-Dihydro-benzo[1,3]dioxo-6-ylmethyl)-amino]-methyl}-piperidin-1-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol




15

The title compound was produced in a manner analogous to Example 1f in a yield of 44% (MS (EI) m/z 450 [M+H]⁺).

Example 10: 4-[(Benzo[1.3]dioxol-5-ylmethyl)-amino]-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol

10.a) 8-([2-(6-Methoxy-quinazolin-4-yl)-ethyl]-1,4-dioxaspiro-[4.5]decan-8-ol




Platinum oxide (0.46 g) was added to a solution of 8-(6-methoxy-quinazolin-4-ylethynyl)-1,4-dioxaspiro[4.5]decan-8-ol (0.96 g, 2.88 mmol) in EtOH (40 ml) and THF (10 ml), and hydrogenation was carried out under a hydrogen atmosphere (1 bar). The reaction mixture was stirred in the presence of activated carbon (5 g) and filtered. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (EtOAc and then EtOAc:MeOH 9/1).

Yield: 0.623 g (1.81 mmol), foam

MS (EI) m/z 344 [M+H]⁺

8-(6-Methoxy-quinazolin-4-ylethynyl)-1,4-dioxaspiro[4.5]decan-8-ol was prepared according to J. Chem. Soc. Perk. Trans. 1, 2000, 3382.

10.b) 4-Hydroxy-4-[2-(6-methoxyquinazolin-4-yl)-ethyl]-cyclo-hexanone




5

A solution of 8-([2-(6-methoxy-quinazolin-4-yl)-ethyl]-1,4-dioxo-spiro[4.5]decan-8-ol (0.623 g, 1.81 mmol) in AcOH/THF/H₂O (3/2/2, 10 ml) was stirred at 60 °C for 30 hours. The reaction mixture was concentrated to dryness by rotary evaporation. The residue was diluted with sodium bicarbonate (100 ml) and ethyl acetate (100 ml), the phases were separated and the aqueous phase was extracted twice using 50 ml of ethyl acetate each time. The combined organic phases were dried over MgSO₄, filtered and concentrated to dryness by rotary evaporation.

Yield: 0.425 g (1.41 mmol) foam.

MS (EI) m/z 301 [M+H]⁺

10.c) 4-[(Benzo[1.3]dioxol-5-ylmethyl)-amino]-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol:




Piperonylamine (0.030 ml, 0.24 mmol) and sodium triacetoxymethylborohydride (0.08 g, 0.377 mmol) were added to a solution of 4-hydroxy-4-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanone (0.06 g, 0.2 mmol) in dichloromethane (1 ml). The reaction mixture was stirred overnight and then filtered over Hydromatrix (wetted with an NaHCO_3 solution, 2 ml) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane:MeOH 9/1 and then dichloromethane:MeOH 9/1 and 2% triethylamine).

Yield: 0.074 g (0.17 mmol) *cis/trans* mixture

MS (EI) m/z 436 $[\text{M}+\text{H}]^+$

15 Example 11: 4-[(Benzo[1,2,5]thiadiazol-5-ylmethyl)-amino]-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol

11.a) 4-Amino-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol



20

Ammonium acetate (14 g, 44 mmol) and sodium triacetoxymethylborohydride (1.2 g, 5 mmol) were added to a solution of 4-hydroxy-4-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanone (1.25 g, 4.2 mmol) in methanol (50 ml). The reaction mixture was stirred overnight and concentrated to dryness by rotary evaporation. The residue was diluted with water (100 ml) and dichloromethane (100 ml), the phases were separated and the aqueous phase was extracted


25

twice using 50 ml of dichloromethane each time. The combined organic phases were dried over Na_2SO_4 , filtered and concentrated to dryness by rotary evaporation.

Yield: 0.85 g. 2.82 mmol (oil) *cis/trans* mixture

5 MS (EI) m/z 302 $[\text{M}+\text{H}]^+$

11.b) 4-[(Benzo[1,2,5]thiadiazol-5-ylmethyl)-amino]-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol




10 Molecular sieve, type 3A, (1 g) and benzo[1,2,5]thiadiazol-5-carbaldehyde (0.066 g, 0.4 mmol) were added to a solution of 4-amino-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol (0.116 g, 0.38 mmol) in methanol (1 ml) and dichloroethane (3 ml). The reaction mixture was stirred at room temperature for 20 hours and sodium triacetoxymethylborohydride (0.22 g, 1.04 mmol) was added. The reaction mixture was stirred at room temperature for 2 hours and then filtered over Hydromatrix (wetted with an NaHCO_3 solution, 2 ml) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 9/1 1% NH_4OH).

Yield: 0.112 g. 0.24 mmol (oil) *cis/trans*-mixture.

25 MS (EI) m/z 450 $[\text{M}+\text{H}]^+$

Example 12: 1-[2-(6-Methoxy-quinazolin-4-yl)-ethyl]-4-(3-phenyl-allylamino)-cyclohexanol




5

In an analogous manner, starting from 4-amino-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol (Example 11.b), 1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-4-(3-phenyl-allylamino)-cyclohexanol in the form of a *cis/trans* mixture was produced in a yield of 54% (MS (EI) m/z 418 $[M+H]^+$).

10

Example 13: 4-(3-Furan-2-yl-allylamino)-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol


15



In an analogous manner, starting from 4-amino-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol (Example 11.b), 4-(3-furan-2-yl-allylamino)-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol in the form of a *cis/trans* mixture was produced in a yield of 62% (MS (EI) m/z 408 $[M+H]^+$).

20

Example 14: 4-[(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-
amino]-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclo-
hexanol




5

In an analogous manner, starting from 4-amino-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol (Example 11.b), 4-[(2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol in the form of a *cis/trans* mixture was produced in a yield of 68% (MS (EI) m/z 450 $[M+H]^+$).

10

Example 15: 1-[2-(6-Methoxy-quinazolin-4-yl)-ethyl]-4-
[(quinoxalin-2-ylmethyl)-amino]-cyclohexanol




15

In an analogous manner, starting from 4-amino-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-cyclohexanol (Example 11.b), 1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]-4-[(quinoxalin-2-ylmethyl)-amino]-cyclohexanol in the form of a *cis/trans* mixture was produced in a yield of 59% (MS (EI) m/z 450 $[M+H]^+$).

20

Example 16: cis- and trans-4-[(Benzo[1,3]dioxol-5-ylmethyl)-amino]-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol

- 5 16 a) 8-(6-Methoxy-[1,5]naphthyridin-4-ylethynyl)-1,4-dioxo-spiro[4.5]decan-8-ol




A degassed solution of trifluoromethanesulphonic acid 6-methoxy-[1,5]naphthyridin-4-yl ester (986 mg, 3.2 mmol)
 10 (WO 03 010138) and 8-ethynyl-1,4-dioxo-spiro[4.5]decan-8-ol (638 mg, 3.5 mmol) (prepared according to J. Chem. Soc. Perk. Trans. 1, 2000, 3382) in DMF (3 ml) was added dropwise to a likewise degassed suspension of copper(I) iodide (50 mg) and PdCl₂(PPh₃)₂ (100 mg) in DMF (2 ml) and
 15 triethylamine (3 ml). The reaction mixture was stirred at room temperature for 30 minutes and diluted with water and ether, and the organic phase was washed with water, saturated ammonium chloride solution and saturated sodium chloride solution, dried over MgSO₄ and concentrated. The
 20 crude product was purified by chromatography on silica gel (hex/EtOAc 1:1, EtOAc).

Yield: 0.965 g. 2.8 mmol (oil).

MS (EI) m/z 341 [M+H]⁺

16.b) 8-[2-(6-Methoxy-[1,5]naphthyridin-4-yl)-ethyl]-1,4-dioxo-spiro[4.5]decan-8-ol




5 A solution of 8-(6-methoxy-[1,5]naphthyridin-4-ylethynyl)-1,4-dioxo-spiro[4.5]decan-8-ol (965 mg, 2.8 mmol) in ethanol (100 ml) was hydrogenated for 6 hours over PtO₂ (200 mg) at 1 bar of hydrogen. The catalyst was filtered off and replaced by fresh catalyst and the hydrogenation
10 was continued for a further 3 hours. The catalyst was filtered off and the solvent was concentrated. The crude product was purified by chromatography on silica gel (EtOAc).

Yield: 0.960 g, 2.79 mol (oil).

15 MS (EI) m/z 334.4 [M+H]⁺

16.c) 4-Hydroxy-4-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanone




20 A solution of 8-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-1,4-dioxo-spiro[4.5]decan-8-ol (960 mg, 2.79 mmol) in THF/H₂O/AcOH (2:2:3, 22ml) was stirred overnight at 65°C. The reaction mixture was concentrated and purified by chromatography on silica gel (EtOAc).

Yield: 0.700 g, 2.33 mmol (oil).

MS (EI) m/z 301 $[M+H]^+$


16.d) *cis*- and *trans*-4-[(Benzo[1,3]dioxol-5-ylmethyl)-
5 amino]-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-
cyclohexanol



NaBH(OAc)₃ (100 mg) was added to a solution of 4-hydroxy-
10 4-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclo-
hexanone (100 mg, 0.3 mmol) and piperonylamine (100 μ l) in
THF (3 ml). The reaction mixture was stirred at room
temperature for 4 hours, and diluted with dichloromethane
and ammonium hydroxide. The organic phase was dried over
15 MgSO₄ and concentrated. The residue was purified by
chromatography on silica gel (EtOAc/MeOH 9:1 +1% NH₄OH).
cis-4-[(Benzo[1,3]dioxol-5-ylmethyl)-amino]-1-[2-(6-
methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol
(38 mg, MS (EI) m/z 357 $[M+H]^+$) and *trans*-4-[(benzo-
20 [1,3]dioxol-5-ylmethyl)-amino]-1-[2-(6-methoxy-[1,5]-
naphthyridin-4-yl)-ethyl]-cyclohexanol (63 mg, MS (EI)
 m/z 357 $[M+H]^+$) were obtained.

Example 17: *cis*- and *trans*-4-[(Benzo[1,3]dioxo-5-yl-methyl)-amino]-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol

- 5 17.a) 4-[2-(6-Ethoxy-quinolin-4-yl)-ethyl]-4-hydroxy-cyclohexanone




- 10 In a manner analogous to Example 16.c, 4-[2-(6-ethoxy-quinolin-4-yl)-ethyl]-4-hydroxy-cyclohexanone was prepared from 6-ethoxy-quinolin-4-ol. MS (EI) m/z 314 $[M+H]^+$

6-Ethoxyquinolin-4-ol has already been described in Synth. Comm. 2002, 32, 3185.

15

17.b) *cis*- and *trans*-4-[(Benzo[1,3]dioxo-5-ylmethyl)-amino]-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol



20


In a manner analogous to Example 16.d, there were obtained *cis*-4-[(benzo[1,3]dioxol-5-ylmethyl)-amino]-1-[2-(6-methoxy[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol (48%

yield, MS (EI) m/z 449 $[M+H]^+$) and *trans*-4-[(benzo[1,3]-dioxol-5-ylmethyl)-amino]-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol (29% yield, MS (EI) m/z 449 $[M+H]^+$).

5

Example 18: Synthesis of *cis*- and *trans*-4-[(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-amino]-1-[2-(6-methoxy-[1,5]-naphthyridin-4-yl)-ethyl]-cyclohexanol

10 18.a) 4-Amino-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol




Sodium triacetoxyborohydride (1.89 g, 8.75 mmol) was added to a solution of 4-hydroxy-4-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanone (3 g, 8.5 mmol) and ammonium acetate (25.5 g) in methanol (65 ml). The reaction mixture was stirred at room temperature for 3 hours, 0.5 g of sodium triacetoxyborohydride was added and stirring was continued for a further 2 hours. The reaction mixture was diluted with NH_4OH and extracted with EtOAc and dichloromethane. Organic phases were dried over $MgSO_4$ and concentrated.

Yield: 2.6 g, (oil), *cis/trans* mixture

MS (EI) m/z 302 $[M+H]^+$

25

18.b) *cis*- and *trans*-4-[(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol




5

Molecular sieve, type 3A, (2.5 g) was added to a solution of 4-amino-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol (1250 mg, 4.14 mmol) and 2,3-dihydro-benzo[1,4]dioxin-6-carbaldehyde (680 mg, 4.14 mmol) in methanol (25 ml) and THF (25 ml). The reaction mixture was stirred at room temperature for 20 hours and sodium borohydride (157 mg, 4.14 mmol) was added. The reaction mixture was stirred at room temperature for 1 hour and then filtered over Hydromatrix (wetted with NH₄OH) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (EtOAc/MeOH 9/1, 1% NH₄OH).

cis-4-[(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol (640 mg, 34%, MS (EI) *m/z* 450 [M+H]⁺) and *trans*-4-[(2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol (360 mg, 19%, MS (EI) *m/z* 450 [M+H]⁺) were obtained.

25

Example 19: *cis*- and *trans*-6-({4-Hydroxy-4-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexylamino}-methyl)-4H-benzo[1,4]oxazin-3-one




5
 Starting from 4-amino-1-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexanol and 3-oxo-3,4-dihydro-2H-benzo[1,4]oxazine-6-carbaldehyde, there were obtained, in a manner analogous to Example 18, *cis*-6-({4-hydroxy-4-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexylamino}-methyl)-4H-benzo[1,4]oxazin-3-one (330 mg, 17%, MS (EI) m/z 464 [M+H]⁺) and *trans*-6-({4-hydroxy-4-[2-(6-methoxy-[1,5]naphthyridin-4-yl)-ethyl]-cyclohexylamino}-methyl)-4H-benzo[1,4]oxazin-3-one (980 mg, 44%, MS (EI) m/z 464 [M+H]⁺).
 15

Example 20: *cis*- and *trans*-4-[(2,3-Dihydro-benzo[1,4]-dioxin-6-ylmethyl)-amino]-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol

20


20.a) 4-Amino-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol



25

4-Amino-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol was obtained analogously to the sequence of Examples 14 and 15 starting from trifluoromethanesulphonic acid 6-methoxy-quinolin-4-yl ester and 8-ethenyl-1,4-dioxaspiro[4.5]decan-8-ol.

20.b) 4-[(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol




10

Starting from 4-amino-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol and 2,3-dihydro-benzo[1,4]dioxin-6-carbaldehyde, there were obtained, in analogy to Example 18, *cis*-4-[(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-amino]-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol (50 mg, MS (EI) m/z 449.5 $[M+H]^+$) and *trans*-4-[(2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol (56 mg, MS (EI) m/z 449.5 $[M+H]^+$).

20

Example 21: *cis*- and *trans*-6-({4-Hydroxy-4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexylamino}-methyl)-4H-benzo[1,4]oxazin-3-one



25


Starting from 4-amino-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol and 3-oxo-3,4-dihydro-2H-benzo[1,4]-oxazine-6-carbaldehyde, there were obtained, in analogy to

5 Example 18, *cis*-6-((4-hydroxy-4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexylamino)-methyl)-4H-benzo[1,4]oxazin-3-one (32 mg, MS (EI) m/z 462.6 $[M+H]^+$) and *trans*-6-((4-hydroxy-4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexylamino)-methyl)-4H-benzo[1,4]oxazin-3-one (40 mg, MS (EI)

10 m/z 462.6 $[M+H]^+$).

Example 22: 4-[(2,3-Dihydro-[1,4]dioxino[2,3-c]pyridin-7-ylmethyl)-amino]-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol

15




In analogy to Example 11.b, 4-[(2,3-dihydro-[1,4]dioxino[2,3-c]pyridin-7-ylmethyl)-amino]-1-[2-(6-methoxyquinolin-4-yl)-ethyl]-cyclohexanol was obtained in

20 a yield of 72% in the form of a *cis/trans* mixture (MS (EI) m/z 450 $[M+H]^+$).

2,3-Dihydro-[1,4]dioxino[2,3-c]pyridine-7-carbaldehyde was prepared according to WO 03/010138.


25

Example 23: 4-[(Benzo[1,2,5]thiadiazol-5-ylmethyl)-amino]-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol




5 In analogy to Example 11.b, 4-[(benzo[1,2,5]thiadiazol-5-ylmethyl)-amino]-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol was obtained in a yield of 47% in the form of a *cis/trans* mixture (MS (EI) m/z 449 $[M+H]^+$).

10 Example 24: 1-[2-(6-Methoxy-quinolin-4-yl)-ethyl]-4-(3-phenyl-allylamino)-cyclohexanol




15 In analogy to Example 11.b, 1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-4-(3-phenyl-allylamino)-cyclohexanol was obtained in a yield of 46% in the form of a *cis/trans* mixture (MS (EI) m/z 418 $[M+H]^+$).

20 Example 25: 1-[2-(6-Methoxy-quinolin-4-yl)-ethyl]-4-[(quinoxalin-2-ylmethyl)-amino]-cyclohexanol



In analogy to Example 11.b, 1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-4-[(quinoxalin-2-ylmethyl)-amino]-cyclohexanol was obtained in a yield of 61% in the form of a *cis/trans* mixture (MS (EI) m/z 443[M+H]⁺).


Example 26: 6-((4-Hydroxy-4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexylamino)-methyl)-2,3-dihydro-benzo[1,4]-dioxin-5-ol



In analogy to Example 11.b, 6-((4-hydroxy-4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexylamino)-methyl)-2,3-dihydro-benzo[1,4]dioxin-5-ol was obtained in a yield of 19% in the form of a *cis/trans* mixture (MS (EI) m/z 465[M+H]⁺).

The corresponding aldehyde was prepared analogously to J. heterocycl. Chem **1989**, 26, 193-197.


Example 27: 4-[(2-Chloro-quinolin-3-ylmethyl)-amino]-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol



In analogy to Example 11.b, 4-[(2-chloro-quinolin-3-yl-methyl)-amino]-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol was obtained in a yield of 56% in the form of a *cis/trans* mixture (MS (EI) m/z 477 $[M+H]^+$).

5

Example 28: Synthesis of 2-[4-hydroxy-4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexylamino]-N-pyridin-2-yl-acetamide



10

2-Bromo-N-pyridin-2-yl-acetamide (0.068 g, 0.31 mmol) and potassium carbonate (0.045 g, 0.34 mmol) were added to a solution of 4-amino-1-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanol (0.094 g, 0.31 mmol) in DMF (3.5 ml). The reaction mixture was stirred at room temperature for 2 days and then concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 19/1 1% ammonium hydroxide).

15

Yield: 0.11g (0.25 mmol) in the form of a *cis/trans* mixture.

20

MS (EI) m/z 435 $[M+H]^+$


2-Bromo-N-pyridin-2-yl-acetamide has already been described in WO 02/24684.

25

Example 29: Benzo[1,3]dioxol-5-ylmethyl-(4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexyl)-amine

29.a) 8-Ethynyl-1,4-dioxaspiro[4.5]decane:

5




A solution of carbon tetrabromide (12.4 g, 37.4 mmol) in dichloromethane (40 ml) was added at -30°C to a solution of triphenylphosphine (19.6 g, 74.6 mmol) and 1,4-dioxaspiro[4.5]dec-8-carbaldehyde (5 g, 29.37 mmol) in dichloromethane (100 ml). After the solution had been stirred at room temperature for 2 hours, it was concentrated to dryness by rotary evaporation. The residue was diluted with ethyl acetate and n-hexane (1:3; 500 ml), filtered over Celite and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (hexane/EtOAc 4/1). Yield: 6.08 g (18.6 mmol). That material was dissolved in THF (90 ml) and, at -78°C, *n*-BuLi (16.5 ml, 38 mmol 2.3N in hexane) was added dropwise. After the solution had been stirred at -78°C for 1 hour, a solution of 10% NaHSO₄ (50 ml) was added. The aqueous phase was extracted three times using 50 ml of ethyl acetate each time, dried over MgSO₄, filtered and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (hexane/EtOAc 5/1).

Yield: 2.74 g (16.5 mmol)

δ H (CDCl₃, 300 MHz): 1.61 (m, 2H); 1.70-1.94 (m, 6H); 2.07 (d, J=2.5 Hz, 1H); 2.51 (m, 1H); 3.96 (s, 4H).

30

29.b) 4-(1,4-Dioxaspiro[4.5]dec-8-ylethynyl)-6-methoxy-quinoline:




5 A degassed solution of trifluoromethanesulphonic acid 6-methoxyquinolin-4-yl ester (0.95 g, 3.1 mmol) and 4-(1,4-dioxaspiro[4.5]dec-8-ylethynyl)-6-methoxyquinoline (0.514 g, 3.1 mmol) in DMF (6 ml) and TEA (12 ml) was added to a mixture of $\text{PdCl}_2(\text{PPh}_3)_2$ (0.110 g, 0.157 mmol) and CuI
10 (0.055 g, 0.288 mmol). After the solution had been stirred at room temperature for 1 hour, it was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (EtOAc).

Yield: 0.83 g (2.56 mmol)

15 MS (EI) m/z 324 $[\text{M}+\text{H}]^+$

29.c) 4-[2-(1,4-Dioxaspiro[4.5]dec-8-yl)ethyl]-6-methoxyquinoline:




20 Platinum oxide ((0.462 g) was added to a solution of 4-(1,4-dioxaspiro[4.5]dec-8-yl-ethynyl)-6-methoxyquinoline (0.83 g, 2.53 mmol) in EtOH (30 ml) and ethyl acetate (10 ml) and hydrogenation was carried out under a hydrogen atmosphere (1bar). The catalyst was filtered off and the
25 filtrate was concentrated to dryness by rotary

evaporation. The residue was purified by column chromatography on silica gel (EtOAc and then EtOAc/-methanol 9/1).

Yield: 0.77 g (2.35 mmol)

5 MS (EI) m/z 328 [M+H]⁺

29.d) 4-[2-(6-Methoxy-quinolin-4-yl)-ethyl]-cyclohexanone:




10 A solution of 4-[2-(1,4-dioxo-spiro[4.5]dec-8-yl)-ethyl]-6-methoxy-quinoline (0.77 g, 2.35 mmol) in AcOH/THF/H₂O (3/2/2, 10 ml) was stirred at 60°C for 10 hours. The reaction mixture was concentrated to dryness by rotary evaporation. The residue was diluted with EtOAc (100 ml),
15 washed with NaHCO₃ (100 ml), dried over MgSO₄, filtered and concentrated to dryness by rotary evaporation.

Yield: 0.631 g (2.23 mmol)

MS (EI) m/z 284 [M+H]⁺

20 29.e) Benzo[1,3]dioxol-5-ylmethyl-{4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexyl}-amine:




Piperonylamine (0.038 ml, 0.3 mmol) and sodium triacetoxyborohydride (0.05 g) were added to a solution of 4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanone (0.05 g, 0.176 mmol) in dichloromethane (0.5 ml). The reaction mixture was stirred overnight and then filtered over Hydromatrix (wetted with an NaHCO₃ solution, 2 ml) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (EtOAc and then EtOAc:MeOH 9/1).

Yield: 0.069 g (0.165 mmol) in the form of a *cis/trans* mixture

MS (EI) *m/z* 419 [M+H]⁺

Example 30: 6-({4-[2-(6-Methoxy-quinolin-4-yl)-ethyl]-cyclohexylamino}-methyl)-4H-benzo[1,4]oxazin-3-one



30.a) 7-[(Benzyl-{4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexyl}-amino)-methyl]-4H-benzo[1,4]oxazin-3-one


Benzylamine (0.135 ml, 1.24 mmol) and sodium triacetoxyborohydride (0.315 g, 1.55 mmol) were added to a solution of 4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexanone (Example 26.e) (0.35 g, 1.23 mmol) in dichloroethane (8 ml). The reaction mixture was stirred for 2 hours. 3-Oxo-3,4-dihydro-2H-benzo[1,4]oxazine-6-carb-

aldehyde (0.230g, 1.3 mmol), THF (4 ml and sodium tri-acetoxyborohydride (0.315 g, 1.55 mmol) was added. The reaction mixture was stirred overnight and then filtered over Hydromatrix (wetted with an NaHCO_3 solution, 10 ml) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 19/1).

Yield: 0.540 g (1.0 mmol)

MS (EI) m/z 536 $[\text{M}+\text{H}]^+$

30.b) 6-((4-[2-(6-Methoxy-quinolin-4-yl)-ethyl]-cyclohexylamino)-methyl)-4H-benzo[1,4]oxazin-3-one



15


20% $\text{Pd}(\text{OH})_2$ on carbon (0.5 g) was added to a solution of 7-[(benzyl-{4-[2-(6-methoxy-quinolin-4-yl)-ethyl]-cyclohexyl}-amino)-methyl]-4H-benzo[1,4]oxazin-3-one (0.535 g, 1 mmol) in THF (15 ml) and MeOH (5 ml) and hydrogenation was carried out under a hydrogen atmosphere (1 bar). The reaction mixture was filtered and the filtrate was concentrated. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 9/1 1% ammonium hydroxide).

Yield: 0.251 g (0.56 mmol) in the form of a *cis/trans* mixture

MS (EI) m/z 446 $[\text{M}+\text{H}]^+$

Example 31: 2-{3-[(2,3-Dihydro-benzo[1,4]dioxin-6-yl-methyl)amino]pyrrolidin-1-yl}-1-(6-methoxyquinolin-4-yl)-ethanol.

- 5 31.a) Benzyl (3RS)-[(2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-pyrrolidine-1-carbamate




- Sodium bicarbonate (7 g) and chloroformic acid benzyl ester (2.8 ml) were added at 0°C to a solution of 3-amino-pyrrolidine dihydrochloride (3.2 g, 20.1 mmol) in water (100 ml) and acetone (150 ml). The reaction mixture was stirred for 10 hours and then concentrated to dryness by rotary evaporation. The residue was diluted with EtOAc (100 ml). The two phases were separated and the aqueous phase was extracted twice using 50 ml of EtOAc each time. The combined organic phases were washed with 20 ml of saturated sodium chloride solution, dried over MgSO₄, filtered and concentrated to dryness by rotary evaporation. The residue was dissolved in dichloromethane (30 ml), and 1,4-benzodioxane-6-carbaldehyde (1.6 g) and, after 20 minutes, sodium triacetoxyborohydride (4 g) were added. The reaction mixture was stirred overnight and then diluted with sodium bicarbonate (80 ml). The two phases were separated and the aqueous phase was extracted twice using 50 ml of dichloromethane each time. The combined organic phases were washed with 20 ml of saturated sodium chloride solution, dried over MgSO₄, filtered and concentrated to dryness by rotary evaporation. The residue

was purified by column chromatography on silica gel (EtOAc:MeOH 9/1).

Yield: 3.75 g (10.1 mmol)

5 δ H (CDCl₃, 300 MHz): 1.57 (br s, 1H), 1.77 (m, 1H), 2.07 (m, 1H); 3.21 (m, 1H); 3.37 (m, 2H); 3.61 (m, 2H), 3.70 (s, 2H); 4.25 (s, 4H); 5.14 (s, 2H); 6.76-6.84 (m, 3H); 7.30-7.38 (m, 5H).

31.b) (2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)pyrrolidin-
10 (3RS)-yl-amine:




20% Pd(OH)₂ on carbon (1 g) was added to a solution of benzyl 3-[(2,3-dihydro-benzo[1,4]dioxin-6-yl-methyl)-
15 amino]pyrrolidin-1-carbamate (3.75 g, 10.1 mmol) in EtOH (20 ml) and EtOAc (20 ml) and hydrogenation was carried out under a hydrogen atmosphere (1 bar). The reaction mixture was filtered and the filtrate was concentrated.

Yield: 2.15 g (9.2 mmol)

20 MS (EI) m/z 235 [M+H]⁺

31.c) 2-((3RS)-((2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino)-pyrrolidin-1-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol:



5


A solution of 6-methoxy-4-oxiranylquinoline (0.201 g, 1 mmol) and (2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-pyrrolidin-3-ylamine (0.234 g, 1 mmol) in ethanol (2 ml) was heated at 80°C for 16 hours. The reaction mixture was cooled to room temperature and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (EtOAc and then EtOAc/MeOH 5:1).

Yield: 0.208 g (0.477 mmol)

MS (EI) m/z 436.5 [M+H]⁺

Example 32: 4-(4-Methoxy-benzylamino)-1-[2-(6-methoxy-quinazolin-4-yl)-ethyl]cyclohexanol.

20




4-Methoxybenzylamine (0.026 ml, 0.24 mmol) and then sodium triacetoxyborohydride (0.08 g, 0.377 mmol) were added to a

solution of 4-hydroxy-4-[2-(6-methoxyquinazolin-4-yl)-ethyl]cyclohexanone (0.06 g, 0.2 mmol) in dichloromethane (1 ml). The reaction mixture was stirred overnight and then filtered over Hydromatrix (wetted with an NaHCO_3 solution, 2 ml) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 9/1 and then dichloromethane/MeOH 9/1 and 2% triethylamine).

10 Yield: 0.052 g (0.124 mmol) in the form of a *cis/trans* mixture

MS (EI) m/z 422 $[\text{M}+\text{H}]^+$

15 Example 33: 4-(3-Fluorobenzylamino)-1-[2-(6-methoxyquinazolin-4-yl)-ethyl]cyclohexanol.




The compound was prepared analogously to Example 10, starting from 3-fluorobenzylamine (0.026 ml, 0.24 mmol).

20 Yield: (0.052 g, 0.124 mmol) in the form of a *cis/trans* mixture

MS (EI) m/z 410 $[\text{M}+\text{H}]^+$

Example 34: 2-{4-[(2,3-Dihydrobenzo[1,4]dioxin-6-yl-ethyl)-amino]-azepan-1-yl}-1-(6-methoxy-quinolin-4-yl)-ethanol.

5 34.a) *tert*-Butyl 4-oxoazepane 1-carbamate:




A mixture of 5-oxoazepane-1,4-dicarboxylic acid 1-*tert*-butyl ester 4-ethyl ester (5.7 g, 20 mmol; prepared as
 10 described in Synthetic Communications 1992, 22, 1249) was refluxed for 3 hours in a mixture of 3N NaOH (50 ml) and THF (25 ml). The reaction mixture was cooled and neutralised with dilute HCl. The mixture was extracted with ethyl acetate and the organic phases were dried over
 15 MgSO₄ and concentrated.

Yield: 4.2 g (100%)

¹H-NMR (CDCl₃, 300 MHz): 1.46 (s, 9H); 1.80 (br, 2H); 2.60-2.7 (m, 4H); 3.4-3.6 (m, 4H).

20 34.b) *tert*-Butyl (4*RS*)-aminoazepane-1-carbamate:



Ammonium acetate (3.5 g) and sodium cyanoborohydride (295 mg, 1 eq.) were added to a solution of *tert*-butyl 4-oxoazepane-1-carbamate (1 g, 4.68 mmol) in methanol
 25 (50 ml). The mixture was stirred overnight at room temperature. The reaction mixture was concentrated and the residue was dissolved in saturated potassium carbonate


solution and ethyl acetate. The aqueous phase was extracted with ethyl acetate (2x50ml), and the combined organic phases were dried over magnesium sulphate and concentrated. 1 g (100%) of product was obtained, which was further used without purification.

$^1\text{H-NMR}$ (CDCl_3 , 300 MHz): 1.46 (s, 9H); 1.5-1.75 (m, 2H); 1.8-2.05 (m, 4H); 2.4 (br, 2H); 2.95-3.05 (m, 1H); 3.1-3.6 (m, 4H).

MS (EI) m/z 215.6 $[\text{M}+\text{H}]^+$

10

34.c) *tert*-Butyl (4*RS*)-[(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-amino]-azepane-1-carbamate:



Acetic acid (500 μl) and sodium cyanoborohydride (126 mg) were added to a solution of *tert*-butyl 4-aminoazepane-1-carbamate (428 mg, 2 mmol) and 2,3-dihydrobenzo[1,4]dioxin-6-carbaldehyde (330 μl , 2 mmol) in dichloroethane (10 ml). The mixture was stirred overnight at room temperature, diluted with saturated potassium carbonate solution and extracted with dichloromethane. The organic phases were dried over magnesium sulphate and concentrated. The product was purified by chromatography on silica gel (ethyl acetate).


Yield: 370 mg (51%)

$^1\text{H-NMR}$ (CDCl_3 , 300 MHz): 1.46 (s, 9H); 1.5-1.75 (m, 2H); 1.8-2.05 (m, 4H); 2.65 (m, 1H); 3.1-3.6 (m, 4H); 3.70 (s, 2H); 4.26 (s, 4H); 6.8-6.9 (m, 3H).

MS (EI) m/z 363.6 $[\text{M}+\text{H}]^+$

30

34.d) Azepan-(4RS)-yl-(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-amine:




5 *tert*-Butyl 4-[(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-amino]-azepane-1-carbamate (370 mg, 1 mmol) was dissolved in a mixture of 10 ml of water and 2 ml of conc. HCl and stirred at room temperature for 24 hours. The reaction mixture was neutralised with solid potassium carbonate and
10 extracted with ethyl acetate. Chromatography on silica gel (dichloromethane/MeOH 9:1) yielded 220 mg of pure product (85%).

¹H-NMR (CDCl₃, 300 MHz): 1.8-2.0 (m, 1H); 2.0-2.3 (m, 4H); 2.3-2.6 (m, 2H); 3.2-3.4 (m, 4H); 3.6-6.7 (m, 1H);
15 3.95 (dd, 2H); 4.2-4.3 (m, 5H); 6.8-7.1 (m, 3H).

MS (EI) m/z 263.4 [M+H]⁺

34.e) 2-[(4RS)-[(2,3-Dihydrobenzo[1,4]dioxin-6-ylmethyl)-amino]-azepan-1-yl]-(1RS)-(6-methoxyquinolin-4-yl)ethanol:



20

A mixture of azepan-4-yl-(2,3-dihydro-benzo[1,4]-dioxin-6-ylmethyl)amine (60 mg), 6-methoxy-4-oxiranyl-quinoline (50 mg), lithium perchlorate (25 mg) and potassium carbonate (35 mg) was heated overnight at 80°C in DMF
25 (1 ml). The reaction mixture was purified by

chromatography on silica gel (dichloromethane/MeOH 9:1 (+2% NEt₃)).


¹H-NMR (CDCl₃, 300 MHz): 1.5-2.0 (m, 7H); 2.45 (dd, 1H); 2.55-3.05 (m, 8H); 3.5-3.8 (m, 2H); 3.85 (s, 3H);
 5 4.15 (s, 4H); 5.25-5.35 (m, 1H); 6.6-6.8 (m, 2H); 7.11 (dd, 1H); 7.30 (dd, 1H); 7.55 (dd, 1H); 7.97 (d, 1H); 8.7 (d, 1H).

MS (EI) m/z 464.6 [M+H]⁺

10 Example 35: [1-(2,3-Dihydro-benzo[1,4]dioxin-6-yl-methyl)-azepan-4-yl] (6-methoxyquinolin-4-ylmethyl)amine:

35.a) *tert*-Butyl (4*RS*)-[(6-methoxyquinolin-4-ylmethyl)-amino]-azepane-1-carbamate:

15




Sodium cyanoborohydride (170 mg, 2.7 mmol) was added to a solution of *tert*-butyl 4-aminoazepane-1-carbamate (568 mg, 2.65 mmol) and 6-methoxyquinoline-4-carbaldehyde (497 mg, 2.7 mmol) in dichloroethane (10 ml) and acetic acid (1 ml). The reaction mixture was stirred at room temperature overnight, poured onto saturated sodium carbonate solution and extracted with dichloromethane. The organic phase was dried over magnesium sulphate and
 20 concentrated. The product was purified by chromatography on silica gel (ethyl acetate).

Yield: 515 mg (51%)

¹H-NMR (CDCl₃, 300 MHz): 1.47 (s, 9H); 1.5-2.0 (m, 8H);
 2.8-2.9 (m, 1H); 3.2-3.65 (m, 5H); 3.97 (s, 3H); 4.22
 (s, 2H); 5.25-5.35 (m, 1H); 6.6-6.8 (m, 2H); 7.11 (dd,
 1H); 7.35-7.45 (m, 3H); 8.02 (d, *J*=9.2Hz, 1H); 8.73 (d,
 5 *J*=1.4Hz, 1H).
 MS (EI) *m/z* 386.5 [M+H]⁺

35.b) Azepan-(4RS)-yl-(6-methoxyquinolin-4-ylmethyl)amine:



10


TFA (1 ml) was added at room temperature to a solution of
tert-butyl 4-[(6-methoxyquinolin-4-ylmethyl)amino]-
 azepane-1-carbamate (700 mg, 1.8 mmol) in dichloromethane
 (1 ml). The reaction mixture was stirred at room
 15 temperature for 3 hours, concentrated and taken up in
 aqueous NaOH. The aqueous phase was extracted with ethyl
 acetate and the combined organic phases were dried over
 sodium sulphate. The product was purified by
 chromatography on silica gel (DCM/MeOH 9:1 (1% NH₄OH)).

20 Yield: 477 mg (92%)

¹H-NMR (CDCl₃, 300 MHz): 1.7-2.3 (m, 6H); 3.0-3.3 (m,
 4H); 3.35-3.45 (m, 1H); 3.97 (s, 3H); 4.2 (s, 2H); 7.28
 (d, 1H); 7.35-7.45 (m, 2H); 8.02 (d, *J*=9.2Hz 1H); 8.73
 (d, *J*=1.4Hz, 1H).

25 MS (EI) *m/z* 286.3 [M+H]⁺

35.c) [1-(2,3-Dihydrobenzo[1,4]dioxin-6-ylmethyl)-azepan-(4RS)-yl]-(6-methoxyquinolin-4-ylmethyl)amine:




5 Sodium triacetoxyborohydride (100 mg, 0.47 mmol) was added to a solution of azepan-4-yl-(6-methoxyquinolin-4-ylmethyl)amine (90 mg, 0.32 mmol) and 2,3-dihydro-benzo-[1,4]dioxin-6-carbaldehyde (51.8 mg, 0.32 mmol) in dichloroethane/THF (1:1, 0.7 ml). The reaction mixture was
10 stirred at room temperature for 3 hours, concentrated and chromatographed on silica gel (ethyl acetate, methanol).

Yield: 56 mg (41%)

¹H-NMR (CDCl₃, 300 MHz): 1.40-2.05 (m, 6H); 2.35-2.80 (m, 3H); 2.8-2.9 (m, 1H); 3.41 (2, 2H); 3.99 (s, 3H);
15 4.15 (s, 2H); 4.20 (s, 4H); 6.70-6.80 (m, 3H); 7.37 (dd, *J*=2.76, *J*=9.1, 1H); 7.45 (d, *J*=2.76, 1H); 7.51 (d, *J*=4.4, 1H); 7.92 (d, *J*=9.12, 1H); 8.65 (d, *J*=4.4, 1H).
MS (EI) *m/z* 434.7 [M+H]⁺


20 Example 36: (6-Methoxyquinolin-4-ylmethyl)-(1-phenethyl)-azepan-(4RS)-yl)-amine:



There was also prepared in a manner analogous to Example 35.c: (6-methoxyquinolin-4-ylmethyl)-(1-phenethylazepan-(4RS)-yl)-amine (25% yield, MS (EI) m/z 390.5 $[M+H]^+$).


5

Example 37: (6-Methoxyquinolin-4-ylmethyl)-[1-(3-phenylpropyl)-azepan-4-yl]amine




10 There was also prepared in a manner analogous to Example 35.c: (6-methoxyquinolin-4-ylmethyl)-[1-(3-phenylpropyl)-azepan-4-yl]amine (25% yield, MS (EI) m/z 404.9 $[M+H]^+$)

15 Example 38: (1-Heptylazepan-(4RS)-yl)-(6-methoxyquinolin-4-ylmethyl)amine:



20 There was also prepared in a manner analogous to Example 35.c: (1-heptylazepan-(4RS)-yl)-(6-methoxyquinolin-4-ylmethyl)amine (43% yield, MS (EI) m/z 384.4 $[M+H]^+$).

Example 39: 1-((4RS)-[(6-methoxyquinolin-4-ylmethyl)-amino]-azepan-1-yl)-3-phenylpropenone.




5 Cinnamic acid chloride (52.5 mg, 1 eq.) was added to a solution of azepan-4-yl-(6-methoxyquinolin-4-yl-methyl)-amine (90 mg, 0.32 mmol) in THF/DCE. The reaction mixture was stirred at room temperature for hours, concentrated, and purified by chromatography on silica gel (DCM/MeOH 9:1
10 (+1% NH₄OH)).

Yield: 37 mg (38%)

MS (EI) m/z 416.6 [M+H]⁺

Example 40: 1-[(2RS)-Hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]-4-phenethyl-[1,4]diazepane-(5RS)-carboxylic acid
15 tert-butyl ester

40.a) 1-[2-Hydroxy-2-(6-methoxyquinolin-4-yl)ethyl]-[1,4]-diazepane-5-carboxylic acid tert-butyl ester




20

A mixture of 6-methoxy-4-oxiranylquinoline (400 mg, 2 mmol), [1,4]-diazepane-5-carboxylic acid tert-butyl

ester (400 mg, 2 mmol, prepared as described in J. Chem. Research (S), 1991, 306, 2876), lithium perchlorate (211 mg, 2 mmol) and potassium carbonate (275 mg, 2 mmol) in DMF (5 ml) was heated at 100°C for 4 hours. The reaction mixture was diluted with water and ethyl acetate, the aqueous phase was extracted with ethyl acetate and the combined organic phases were dried over sodium sulphate and concentrated. The product was purified by chromatography on silica gel (DCM/MeOH 9:1).

Yield: 330 mg (41%)
MS (EI) m/z 402.5 [M+H]⁺

40.b) 1-[(2RS)-Hydroxy-2-(6-methoxyquinolin-4-yl)ethyl]-4-phenethyl-[1,4]diazepane-(5RS)-carboxylic acid *tert*-butyl ester:




Sodium triacetoxyborohydride (79 mg (1.5 eq) was added to a solution of 1-(2-hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl)-[1,4]diazepane-5-carboxylic acid *tert*-butyl ester (100 mg, 0.25 mmol) and phenyl acetaldehyde (29.2 μ l, 1 eq.) in THF (700 μ l). After 2 hours, a further equivalent of phenyl acetaldehyde was added and the reaction mixture was heated to 40°C. After a further 2 hours, the reaction mixture was concentrated to dryness and purified by chromatography on silica gel (ethyl acetate, methanol).

Yield: 67 mg (54%)

MS (EI) m/z 506 [M+H]⁺

- 40.c) 1-[(2RS)-Hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]-4-phenethyl-[1,4]-diazepane-(5RS)-carboxylic acid:



- 1-[(2-Hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]-4-phenethyl-[1,4]diazepane-5-carboxylic acid tert-butyl ester (55 mg) was suspended in 4M HCl in dioxane and stirred at room temperature for 3.5 hours. The reaction mixture was concentrated and purified by preparative HPLC.

Yield: 3.9 mg

MS (EI) m/z 450 [M+H]⁺


15

The following were also synthesised in analogous manner:

- 40.d) 1-[2-hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]-4-(3-phenylpropyl)-[1,4]diazepane-5-carboxylic acid tert-butyl ester and
- 20 40.e) 1-[2-hydroxy-2-(6-methoxyquinolin-4-yl)ethyl]-4-(3-phenylpropyl)-[1,4]diazepane-5-carboxylic acid starting from dihydrocinnamaldehyde.
- In the same way,
- 40.f) 4-heptyl-1-[2-hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]-[1,4]diazepane-5-carboxylic acid tert-butyl ester
- 25 and
- 40.g) 4-heptyl-1-[2-hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]-[1,4]diazepane-5-carboxylic acid

were prepared starting from heptanal.

Example 41: 2-[(RS)5-Hydroxymethyl-4-(3-phenylpropyl)-
[1,4]diazepan-1-yl]- (1RS) - (6-methoxyquinolin-4-yl) -
 5 ethanol.




Lithium aluminium hydride (28 mg) was added to a solution of 1-[2-hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]-4-(3-phenylpropyl)-[1,4]-diazepane-5-carboxylic acid *tert*-butyl ester (80 mg) in THF and stirring was carried out at room temperature for 1 hour. A few drops of saturated Rochelle salt solution were added to the reaction mixture, stirring was carried out for 15 minutes and the precipitate was filtered off.

15 Yield: 39 mg (57%)
 MS (EI) m/z 450 $[M+H]^+$

Starting from 4-heptyl-1-[2-hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]-[1,4]diazepane-5-carboxylic acid *tert*-butyl ester, 2-(4-heptyl-5-hydroxy-methyl-[1,4]-diazepan-1-yl)-1-(6-methoxyquinolin-4-yl)-ethanol was produced in an analogous manner in a yield of 33% (MS (EI) m/z 430 $[M+H]^+$).

Example 42: (2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-[4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexyl]-amine

42.a) 4-(6-Methoxy-quinazolin-4-yloxymethyl)-cyclohexanol




5

Sodium hydride (79 mg (1.5 eq.) was added at 0°C to a solution of 4-hydroxymethyl-cyclohexanol (3 g, 12 mmol, synthesised analogously to J.Org.Chem. **1994** 59 p. 2748-2761) in DMF (30 ml). After 20 minutes, a solution of
 10 4-chloro-6-methoxy-quinazoline (1.94 g, 10 mmol) in DMF (10 ml) was added and the reaction mixture was heated to 40°C. After a further 2 hours, the reaction mixture was concentrated to dryness. The residue was diluted with water (100 ml) and ethyl acetate (100 ml), the phases were
 15 separated and the aqueous phase was extracted twice using 50 ml of ethyl acetate each time. The combined organic phases were dried over MgSO₄, filtered and concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (EtOAc/hexane 1/2).

20 Yield: 2.01 g, 7 mmol (54%)

MS (EI) m/z 289 [M+H]⁺

42.b) 4-(6-Methoxy-quinazolin-4-yloxymethyl)-cyclohexanone




Triethylamine (10 ml, 71 mmol) and then, in portions, Pyr.SO₃ (5.8 g, 36 mmol), were added at 0°C to a solution of 4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexanol (2.88 g, 10 mmol) in DMSO (30 ml). The reaction mixture
5 was stirred at that temperature for 15 minutes and then allowed to come to room temperature. After 2 hours, water (300 ml) was added. The aqueous phase was extracted with ether (3 x 150 ml). The combined organic phases were washed with saturated sodium chloride solution, dried over
10 MgSO₄ and concentrated. The crude product was purified by chromatography on silica gel (EtOAc/hex 1/2).

Yield: 2.6 g, 9.1 mmol

MS (EI) m/z 287.1 [M+H]⁺

15 42.c) Benzyl-[4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexyl]-amine




Benzylamine (1.08 ml, 10 mmol) and sodium cyanoborohydride (2.2 g, 11 mmol) were added to a solution of 4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexanone (2.87 g, 10 mmol)
20 in methanol (50 ml). The mixture was stirred overnight at room temperature. The reaction mixture was concentrated and the residue was dissolved in saturated potassium carbonate solution and ethyl acetate. The aqueous phase
25 was extracted with ethyl acetate (2x50ml), and the combined organic phases were dried over magnesium sulphate and concentrated. The residue was purified by column

chromatography on silica gel (EtOAc and then EtOAc/MeOH 9/1).

Yield: 2.87 g, 10 mmol in the form of a *cis/trans* mixture

5 MS (EI) m/z 287.1 $[M+H]^+$

42.d) 4-(6-Methoxy-quinazolin-4-yloxymethyl)-cyclohexyl-amine




10 20% Pd(OH)₂ on carbon (1 g) was added to a solution of benzyl-[4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexyl]-amine (2 g, 5.3 mmol) in MeOH (30 ml) and hydrogenation was carried out under a hydrogen atmosphere (1 bar) at 65°C. The reaction mixture was filtered and the
15 filtrate was concentrated.

Yield: 1.44 g, 5 mmol in the form of a *cis/trans* mixture

MS (EI) m/z 288 $[M+H]^+$

20 42.e) Title compound




2,3-Dihydro-benzo[1,4]dioxin-6-carbaldehyde (0.045 g, 0.27 mmol) and then sodium triacetoxyborohydride (0.08 g,

0.377 mmol) were added to a solution of 4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexylamine (0.07 g, 0.25 mmol) in dichloroethane (1 ml). The reaction mixture was stirred for 2 hours and then filtered over Hydromatrix (wetted with NaHCO₃ solution, 2 ml) and subsequently washed with dichloromethane. The filtrate was concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 9/1 and then dichloromethane/MeOH 6/1).

Yield: 0.02g (0.046 mmol) in the form of a *cis/trans* mixture


MS (EI) m/z 436 [M+H]⁺

Example 43: [4-(6-Methoxy-quinazolin-4-yloxymethyl)-cyclohexyl]-quinoxalin-2-ylmethyl-amine



In analogy to Example 42.e, [4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexyl]-quinoxalin-2-ylmethylamine was produced in a yield of 31% in the form of a *cis/trans* mixture (MS (EI) m/z 430 [M+H]⁺).

Example 44: 6-([4-(6-Methoxy-quinazolin-4-yloxymethyl)-cyclohexylamino]-methyl)-4H-benzo[1,4]oxazin-3-one




5

In analogy to Example 42.e, 6-([4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexylamino]-methyl)-4H-benzo[1,4]oxazin-3-one was produced in a yield of 38% in the form of a *cis/trans* mixture (MS (EI) m/z 448 $[M+H]^+$).


10

Example 45: [4-(6-Methoxy-quinazolin-4-yloxymethyl)-cyclohexyl]-(3-phenyl-allyl)-amine



15 In analogy to Example 42.e, [4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexyl]-(3-phenyl-allyl)-amine was produced in a yield of 25% in the form of a *cis/trans* mixture (MS (EI) m/z 405 $[M+H]^+$).

Example 46: 2-[4-(6-Methoxy-quinazolin-4-yloxymethyl)-cyclohexylamino]-N-pyridin-2-yl-acetamide




5 2-Bromo-N-pyridin-2-yl-acetamide (0.075 g, 0.35 mmol) and potassium carbonate (0.054 g, 0.39 mmol) were added to a solution of 4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexylamine (0.1 g, 0.35 mmol) in DMF (3.5 ml). The reaction mixture was stirred at room temperature for
10 2 days and then concentrated to dryness by rotary evaporation. The residue was purified by column chromatography on silica gel (dichloromethane/MeOH 19/1 1% NH₄OH).

Yield: 0.086 g (0.2 mmol) in the form of a *cis/trans* mixture

15 MS (EI) *m/z* 422 [M+H]⁺

Example 47: (2,3-Dihydro-[1,4]dioxino[2,3-*c*]pyridin-7-ylmethyl)-[4-(6-methoxy-quinazolin-4-yloxymethyl)-cyclohexyl]-amine




20

Starting from Example 42.e, was produced in an analogous manner in a yield of 55% in the form of a *cis/trans* mixture (MS (EI) *m/z* 437 [M+H]⁺).

Example 48: (2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-[4-(6-methoxy-quinolin-4-yloxymethyl)-cyclohexyl]-amine

48.a) *trans*-4-*tert*-Butoxycarbonylamino-cyclohexane-
5 carboxylic acid




Di-*tert*-butyl dicarbonate (12 g, 55 mmol) dissolved in dioxane (50 ml) was added dropwise to a solution of *trans*-4-aminocyclohexanecarboxylic acid (Synth. Commun. 2002,
10 32, 1985) (7.16 g, 50 mmol) in dioxane (50 ml) and water (50 ml). After stirring for 30 minutes at room temperature, 1M NaOH (50 ml) was added and the reaction mixture was stirred overnight at room temperature. The dioxane was distilled off using a rotary evaporator and
15 the aqueous residue was adjusted to pH 3 using 1N HCl and extracted with ethyl acetate. Organic extracts dried over Na₂SO₄ and concentrated.

Yield: 13.2 g

MS (EI) *m/z* 242.4[M-H]⁻.

20

48.b) *tert*-Butyl *trans*-(4-hydroxymethyl-cyclohexyl)-
carbamate




A solution of borane dimethyl sulphide complex (2.85 ml, 30 mmol) in THF (50 ml) was added dropwise at 0°C to a solution of *trans*-4-*tert*-butoxycarbonylamino-cyclohexane-carboxylic acid (2.43 g, 10 mmol) in THF (100 ml). The
25

reaction mixture was stirred at 0°C for 15 minutes and at room temperature for 3 hours, methanol was cautiously added and the mixture was concentrated. The crude product was repeatedly treated with methanol and concentrated again, and finally dried under a high vacuum. Used without further purification.


Yield: 2.4 g (quant.)

48.c) *trans*-Methanesulphonic acid 4-*tert*-butoxycarbonyl-aminocyclohexylmethyl ester



At 0°C, triethylamine (417 μ l, 2.2 mmol) was added to a solution of *tert*-butyl *trans*-(4-hydroxymethylcyclohexyl)-carbamate (459 mg, 3 mmol) in dichloromethane (25 ml), and methanesulphonyl chloride (171 μ l, 2.2 mmol) was added dropwise. The reaction mixture was stirred at 0°C for 5 hours and at room temperature for 1 hour, taken up with water and extracted with dichloromethane. The organic phase was dried over Na₂SO₄ and concentrated. The crude product was reacted without further purification.


48.d) *tert*-Butyl [4-(6-methoxy-quinolin-4-yloxymethyl)-cyclohexyl]carbamate



NaH dispersion (55% in mineral oil, 88 mg, 2 mmol) was added to a suspension of 6-methoxy-4-hydroxyquinoline (350 mg, 2 mmol) in DMF (4 ml) and stirred at room temperature until a solution formed. *trans*-Methanesulphonic acid 4-*tert*-butoxycarbonylaminocyclohexylmethyl ester (635 mg, 2 mmol) was added and the reaction mixture was stirred overnight at 80°C. Water was added to the mixture and extraction was carried out with ethyl acetate. Organic extracts washed with water, dried over Na₂SO₄ and concentrated. The crude product was purified by chromatography on silica gel (DCM/MeOH 9:1).

Yield: 438 mg, 56.6% (oil)
(MS (EI) *m/z* 387 [M+H]⁺).

48.e) *trans*-4-(6-Methoxy-quinolin-4-yloxymethyl)-cyclohexylamine



A solution of *tert*-butyl [4-(6-methoxy-quinolin-4-yloxymethyl)-cyclohexyl]carbamate (438 mg, 1.13 mmol) and TFA (1.5 ml) in dichloromethane (4 ml) was stirred at room temperature for 2 hours. The reaction mixture was poured


79

onto ice/ammonium hydroxide and extracted with dichloromethane. The organic extracts were dried over Na_2SO_4 and concentrated.

Yield: 313 mg, 97%

5 (MS (EI) m/z 287 $[\text{M}+\text{H}]^+$).

48.f) Title compound




10

In analogy to Example 42.e, *trans*-2,3-dihydro-benzo[1,4]-dioxin-6-ylmethyl)-[4-(6-methoxy-quinolin-4-yloxymethyl)-cyclohexyl]-amine was produced in a yield of 73% (MS (EI) m/z 435 $[\text{M}+\text{H}]^+$).

15


Example 49: 6-([4-(6-Methoxy-quinolin-4-yloxymethyl)-cyclohexylamino]-methyl)-4H-benzo[1,4]oxazin-3-one



20 In analogy to Example 42.e, *trans*-6-([4-(6-methoxy-quinolin-4-yloxymethyl)-cyclohexylamino]-methyl)-4H-benzo[1,4]oxazin-3-one was produced in a yield of 81% (MS (EI) m/z 448 $[\text{M}+\text{H}]^+$).


Example 50: (2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-[4-(6-methoxy-[1,5]naphthyridin-4-yloxymethyl)-cyclohexyl]-amine

- 5 50.a) 4-(6-Methoxy-[1,5]naphthyridin-4-yloxymethyl)-cyclohexylamine




- 4-(6-Methoxy-[1,5]naphthyridin-4-yloxymethyl)cyclohexylamine was prepared analogously to Example 48 starting from
 10 6-methoxy-[1,5]naphthyridin-4-ol.

50.b) Title compound



- In analogy to Example 42.e, (2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-[4-(6-methoxy-[1,5]naphthyridin-4-yloxy-
 15 methyl)-cyclohexyl]-amine was produced in a yield of 55% (MS (EI) m/z 456.6 [M+H]⁺).


Example 51: 6-([4-(6-Methoxy-[1,5]naphthyridin-4-yloxy-methyl)-cyclohexylamino]-methyl)-4H-benzo[1,4]oxazin-3-one



5 In analogy to Example 42.e, *trans*-6-([4-(6-methoxy-[1,5]naphthyridin-4-yloxymethyl)-cyclohexylamino]-methyl)-4H-benzo[1,4]oxazin-3-one was produced in a yield of 79% (MS (EI) m/z 456.6 $[M+H]^+$).

10 Example 52: 2-{3-[(2,3-Dihydro-benzo[1,4]dioxin-6-yl-methyl)-amino]-8-aza-bicyclo[3.2.1]oct-8-yl}-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol


52.a) *tert*-Butyl {8-[(2RS)-hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-8-aza-bicyclo[3.2.1]oct-3-yl}carbamate



In analogy to Example 2.c, the title compound was produced in a yield of 82% (MS (EI) m/z 428 $[M+H]^+$).

20 *tert*-Butyl {8-aza-bicyclo[3.2.1]oct-3-yl}carbamate was prepared according to *Eur. J. Med. Chem.* **1991** (34) p. 646-653.

52.b) 2-(3-[(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-8-aza-bicyclo[3.2.1]oct-8-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol




5

In analogy to Example 2.c, the title compound was produced in a yield of 86% (MS (EI) m/z 328 $[M+H]^+$).

Example 53: 2-(3-[(Benzo[1,3]dioxol-5-ylmethyl)-amino]-8-aza-bicyclo[3.2.1]oct-8-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol


10



In analogy to Example 2.e, 2-(3-[(benzo[1,3]dioxol-5-ylmethyl)-amino]-8-aza-bicyclo[3.2.1]oct-8-yl)-1-(6-methoxy-quinolin-4-yl)-ethanol was produced in a yield of 62% (MS (EI) m/z 462 $[M+H]^+$).

15

Example 54: 6-((8-[(2RS)-Hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-8-aza-bicyclo[3.2.1]oct-3-ylamino)-methyl)-4H-benzo[1,4]oxazin-3-one




5

In analogy to Example 2.e, 6-((8-[2-hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-8-aza-bicyclo[3.2.1]oct-3-ylamino)-methyl)-4H-benzo[1,4]oxazin-3-one was produced in a yield of 7% (MS (EI) m/z 489 $[M+H]^+$).

10

Example 55: 2-{3-[(Benzo[1,2,5]thiadiazol-5-ylmethyl)-amino]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-(6-methoxy-quinolin-4-yl)-ethanol




15

In analogy to Example 2.e, 2-{3-[(benzo[1,2,5]thiadiazol-5-ylmethyl)-amino]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-(6-methoxy-quinolin-4-yl)-ethanol was produced in a yield of 59% (MS (EI) m/z 476 $[M+H]^+$).


20

Example 56: 1-(6-Methoxy-quinolin-4-yl)-2-[3-(3-phenyl-allylamino)-8-aza-bicyclo[3.2.1]oct-8-yl]-ethanol




5 In analogy to Example 2.e, 1-(6-methoxy-quinolin-4-yl)-2-[3-(3-phenyl-allylamino)-8-aza-bicyclo[3.2.1]oct-8-yl]-ethanol was produced in a yield of 44% (MS (EI) m/z 445 $[M+H]^+$).

10 Example 57: 2-[3-(3-Furan-2-yl-allylamino)-8-aza-bicyclo[3.2.1]oct-8-yl]-1-(6-methoxy-quinolin-4-yl)-ethanol



In analogy to Example 2.e, 2-[3-(3-furan-2-yl-allylamino)-
15 8-aza-bicyclo[3.2.1]oct-8-yl]-1-(6-methoxy-quinolin-4-yl)-ethanol was produced in a yield of 51% (MS (EI) m/z 434 $[M+H]^+$).


Example 58: 2-{3-[(Benzofuran-2-ylmethyl)-amino]-8-aza-bicyclo[3.2.1]oct-8-yl}-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol



5 In analogy to Example 2.e, 2-{3-[(benzofuran-2-ylmethyl)-amino]-8-aza-bicyclo[3.2.1]oct-8-yl}-1-(6-methoxy-quinolin-4-yl)-ethanol was produced in a yield of 62% (MS (EI) m/z 458 $[M+H]^+$).

10 Example 59: 2-(5-{[(2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-methyl}-3,6-dihydro-2H-pyridin-1-yl)-1-(6-methoxy-quinolin-4-yl)-ethanol

59.a) *tert*-Butyl 5-azidomethyl-3,6-dihydro-2H-pyridine-1-
15 carbamate




At 0°C, triethylamine (3.7 ml, 26.2 mmol) and, dropwise, methanesulphonyl chloride (1.2 ml, 15.44 mmol) were added
20 to a solution of *tert*-butyl 5-hydroxymethyl-3,6-dihydro-2H-pyridine-1-carbamate (*Tetrahedron* **1998**, 54, 7045-7056, 2.8 g, 13.1 mmol) in DCM (50 ml). The reaction mixture was stirred at 0°C for 20 minutes before a saturated solution of NaHCO₃ (40 ml) was added, and the two phases were

separated. The aqueous phase was extracted with dichloro-
methane (40 ml). The combined organic phases were washed
with saturated sodium chloride solution and dried over
MgSO₄ and concentrated. The crude product was dissolved in
5 DMF (50 ml), and sodium azide (1.7 g, 26.1 mmol) was
added. The mixture was heated to 80°C. Reaction monitored
by LCMS. Once the reaction was complete, the mixture was
cooled and concentrated. The residue was taken up in EtOAc
and water and the aqueous phase was extracted with EtOAc
10 (2 x 100 ml). The combined organic phases were washed with
saturated sodium chloride solution, dried over MgSO₄ and
concentrated. The residue was purified by chromatography
on silica gel (EtOAc/hex 1/6).

Yield: (2.2 g, 9.23 mmol) Oil.

15 MS (EI) m/z 239.4 [M+H]⁺.

59.b) *tert*-Butyl 5-aminomethyl-3,6-dihydro-2H-pyridine-1-
carbamate




20 Triphenylphosphine on polystyrene (5.24 g) was added to a
solution of *tert*-butyl 5-azidomethyl-3,6-dihydro-2H-
pyridine-1-carbamate (1 g, 4.2 mmol) in THF (25 ml) and
water (0.250 ml). The reaction mixture was shaken at room
temperature for 36 hours. The polymer was filtered off and
25 then washed with THF. The filtrate was concentrated.

Yield: (0.893 g, 4.2 mmol)

MS (EI) m/z 213.4 [M+H]⁺.

59.c) *tert*-Butyl 5-[[(2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-methyl]-3,6-dihydro-2H-pyridine-1-carbamate




5 A solution of *tert*-butyl 5-aminomethyl-3,6-dihydro-2H-pyridine-1-carbamate (0.7 g, 3.3 mmol) and 1,4-benzodioxane-6-carbaldehyde (0.54 g, 3.3 mmol) in THF (2 ml) and DCE (4 ml) was stirred at room temperature for 1 hour. Sodium triacetoxyborohydride (0.770 g, 3.81 mmol) was then
 10 added thereto. The reaction mixture was stirred at room temperature for 5 hours and filtered through Hydromatrix (wetted with saturated NaHCO₃ solution) and the filtrate was concentrated. The residue was purified by chromatography on silica gel (EtOAc/hex 2/1 and then
 15 EtOAc/MeOH 9/1).

Yield: 0.16 g, 0.44 mmol (oil).

MS (EI) *m/z* 361 [M+H]⁺.

59.d) (2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-(1,2,5,6-tetrahydropyridin-3-ylmethyl)-amine
 20




A solution of *tert*-butyl 5-[[(2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-methyl]-3,6-dihydro-2H-pyridine-1-carbamate (0.16 g, 0.44 mmol) in TFA (3 ml) was stirred
 25 at room temperature for 30 minutes, concentrated and taken up in saturated bicarbonate solution (30 ml) and

dichloromethane (30 ml). The aqueous phase was extracted with dichloromethane (2 x 30 ml) and the combined organic phases were washed with saturated sodium chloride solution, dried over MgSO_4 and concentrated.

5 Yield: 0.06 g, 0.23 mmol, 52% (oil)

MS (EI) m/z 261 $[\text{M}+\text{H}]^+$.

59.e) Title compound




10 A solution of (2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-
(1,2,5,6-tetrahydro-pyridin-3-ylmethyl)-amine (0.06 g,
0.23 mmol) and 6-methoxy-4-oxiranylquinoline (0.06 g,
0.29 mmol) in EtOH (2 ml) was heated at 80°C for 14 hours.
The reaction mixture was concentrated and the residue was
15 purified by chromatography on silica gel (DCM/MeOH 9/1).

Yield: 0.025 g, 0.054 mmol, 23% (foam)

MS (EI) m/z 462 $[\text{M}+\text{H}]^+$.

Example 60: 2-((2RS)-{[(Benzo[1,3]dioxol-5-ylmethyl)-amino]-methyl}-morpholin-4-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol

5 60.a) (2RS)-Azidomethyl-4-benzyl-morpholine




At 0°C, triethylamine (4.88 g, 48.25 mmol) and, dropwise, methansulphonyl chloride (2.25 ml, 28.95 mmol) were added to a solution of N-benzyl-2-hydroxymethylmorpholine
 10 (*Synthetic Communications*, **1980**, 10(1), 59-73, 5 g, 24.12 mmol) in DCM (60 ml). After 30 minutes, the reaction was terminated by addition of saturated bicarbonate solution (50 ml). The organic phase was washed with saturated sodium chloride solution (50 ml), dried over
 15 MgSO₄ and concentrated.

The crude product was dissolved in DMF (50 ml), and sodium azide (3.11 g, 47.8 mmol) was added. The mixture was heated overnight at 80°C. The mixture was cooled and concentrated. The residue was taken up in ether (100 ml)
 20 and water (50 ml) and the aqueous phase was extracted with ether (2 x 100 ml). The combined organic phases were washed with saturated sodium chloride solution, dried over MgSO₄ and concentrated. The residue was purified by chromatography on silica gel (EtOAc).

25 Yield: (4.23 g, 18 mmol, 75%) Oil.

MS (EI) m/z 236.2 [M+H]⁺.

90




60.b) 2-(4-Benzyl-morpholin-(2RS)-yl)-methanamine

5 A solution of 2-Azidomethyl-4-benzyl-morpholine (4.23 g, 18 mmol) and triphenylphosphine (9.47 g, 36 mmol) in THF/water (10/1, 100 ml) was heated overnight at 60°C. The reaction mixture was concentrated and the residue was taken up in 3N HCl (200 ml) and EtOAc (200 ml). The
 10 aqueous phase was extracted with EtOAc (4*). The aqueous phase was adjusted to pH 12 using NaOH and extracted with EtOAc (2 x 200 ml), dried over MgSO₄ and concentrated.

Yield: (3.64 g, 16.8 mmol, 93%) Oil.

MS (EI) m/z 207.2 [M+H]⁺.

15

60.c) *tert*-Butyl (4-benzyl-morpholin-(2RS)-ylmethyl)-carbamate


At 0°C, triethylamine (1.78 g, 17.74 mmol) and di-*tert*-
 20 butyl dicarbonate (2.32 g, 10.65 mmol) were added to a solution of 2-(4-benzyl-morpholin-2-yl)-methanamine

(1.83 g, 8.87 mmol) in DCM (45 ml). The reaction mixture was stirred at that temperature for 30 minutes and concentrated. The residue was purified by chromatography on silica gel (EtOAc/hex 1:1).

5 Yield: (1.51 g, 4.93 mmol, 51%) Oil.

MS (EI) m/z 307.3 $[M+H]^+$.

60.d) *tert*-Butyl {4-[(2*RS*)-hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-morpholin-2-ylmethyl}carbamate



10

20% Pd(OH)₂ (0.7 g) was added to a solution of *tert*-butyl {4-benzyl-morpholin-2-ylmethyl}carbamate (1.51 g,

4.93 mmol) in THF:MeOH (1/1, 28 ml). The reaction mixture was stirred for 2 hours under a hydrogen atmosphere. The

15 mixture was filtered and the filtrate was concentrated.


The intermediate was dissolved in DMF (15 ml), and 6-methoxy-4-oxiranylquinoline (0.9 g, 4.48 mmol), lithium perchlorate (0.477 g, 4.48 mmol) and potassium carbonate (0.743 g, 5.376 mmol) were added. The mixture was heated

20 at 80°C for 23 hours. After cooling, filtration was carried out and the filtrate was concentrated. The residue was dissolved in EtOAc and washed with water, dried over MgSO₄ and concentrated. The residue was purified by chromatography on silica gel (DCM:MeOH 9:1).

Yield: (1 g, 2.4 mmol, 48%) Oil.

MS (EI) m/z 418.5 $[M+H]^+$.

60.e) 2-((2RS)-Aminomethyl-morpholin-4-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol




TFA (2.5 ml) was added at 0°C to a solution of *tert*-butyl {4-[2-hydroxy-2-(6-methoxy-quinolin-4-yl)-ethyl]-morpholin-2-ylmethyl}carbamate (0.95 g, 2.27 mmol) in DCM (4 ml). The reaction mixture was stirred at room temperature for 1.5 hours and concentrated. The residue was taken up in DCM/MeOH (9/1, 30 ml) and ammonium hydroxide (20 ml). The aqueous phase was extracted with DCM/MeOH (2 x 30 ml) and the combined organic phases were dried over Na₂SO₄ and concentrated.

Yield: (0.718 g, 2.25 mmol, 99%) Oil.

MS (EI) m/z 318.5 $[M+H]^+$.

60.f) 2-(2-(((Benzo[1,3]dioxol-5-ylmethyl)-amino)-methyl)-morpholin-4-yl)-1-(6-methoxy-quinolin-4-yl)-ethanol




Activated 3A molecular sieve (3.485 g) and piperonal
 5 (0.172 g, 1.15 mmol) were added to a solution of 2-(2-aminomethyl-morpholin-4-yl)-1-(6-methoxy-quinolin-4-yl)-ethanol (0.365 g, 1.15 mmol) in DCM (10.5 ml) and methanol (3.5 ml). The mixture was stirred overnight at room temperature before sodium borohydride (0.112 g, 2.9 mmol)
 10 was added. The reaction mixture was stirred at room temperature for a further 2 hours, filtered over Hydromatrix (wetted with NaHCO₃) and concentrated. The residue was purified by chromatography on silica gel (DCM:MeOH 9:1+ 1% NH₄OH).

15 Yield: (0.261 g, 0.57 mmol, 50%) Oil.

MS (EI) m/z 452.5 [M+H]⁺.

Example 61: 2-((2RS)-(((2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino)-methyl)-morpholin-4-yl)-(1RS)-(6-methoxy-quinolin-4-yl)-ethanol
 20



Analogously to Example 60.f, also 2-(2-(((2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino)-methyl)-morpholin-4-

yl)-1-(6-methoxy-quinolin-4-yl)-ethanol was produced in a yield of 33% (MS (EI) m/z 466 [M+H]⁺.)

Anti-bacterial activity:

5

The MHK (μg/ml) of those compounds was measured in relation to the following bacterial strains: *S. aureus* ATCC 29213, *S. aureus* I6, *E. faecalis* ATCC 29212, *E. faecium* vanA E25-1, *H. influenzae* 11, *E. coli* ATCC 25922, *M. catarrhalis* 117, *S. pneumoniae* ATCC 49619.


10 Examples 5-6, 8, 11-16, 18-26, 30, 44-45, 47-58 have an MHK≤0.125 in relation to at least one of the listed strains.

15 Examples 1-4, 7, 9-10, 17, 27-29, 34, 42, 43, 46, 59, 61 have a MHK≤0.5 in relation to at least one of the listed strains.

Patent Claims

1. Compounds of formula (I):

5



wherein


10 A is an oxygen, sulphur or nitrogen atom or a C₁₋₄alkylene, C₂₋₄alkenylene, C₂₋₄alkynylene or C₁₋₄heteroalkylene group,


15 X₁, X₂, X₃, X₄ and X₅ are each independently of the others nitrogen atoms or groups of formula CR²,

R¹ is a hydrogen atom, a halogen atom, a hydroxy group, an alkyloxy group or a heteroalkyloxy group,

20 R² is a hydrogen atom, a halogen atom, or a hydroxy, alkyl, alkenyl, alkynyl or heteroalkyl group,

R³ is selected from the following groups:





5 the radicals R⁴, each independently of any other(s),
are a hydroxy group, a C₁₋₆alkyl group or a
C₁₋₈heteroalkyl group,

10 R⁵ is an alkyl, alkenyl, alkynyl, heteroalkyl, aryl,
heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkyl-
cycloalkyl, heterocycloalkyl, aralkyl or hetero-
aralkyl radical,

15 n is 0, 1, 2 or 3 and

15

m is 0 or 2,

20 or a pharmacologically acceptable salt, solvate,
hydrate or a pharmacologically acceptable formulation
thereof.

2. Compounds according to claim 1, wherein A is an
oxygen atom or a group of formula CH₂ or CH(OH).


25 3. Compounds according to claim 1 or 2, wherein one of
the groups X₁, X₂, X₃, X₄ and X₅ is a nitrogen atom

and the others are CH groups, or all of the groups X_1 , X_2 , X_3 , X_4 and X_5 are CH groups.

4. Compounds according to any one of claims 1 to 3,
5 wherein R^1 is a halogen atom, a C_{1-6} alkyloxy group, a methyl group or an ethyl group.
5. Compounds according to any one of claims 1 to 4,
10 wherein R^1 is a methoxy group.
6. Compounds according to any one of claims 1 to 5,
wherein R^4 is a C_{1-6} heteroalkyl group having one or two oxygen atoms as individual hetero atoms.
- 15 7. Compounds according to any one of claims 1 to 6,
wherein R^4 is a group of formula $-COOH$, $-CH_2COOH$, $-CH_2CH_2COOH$, $-CH_2COOCH_3$, $-CH_2CH_3$, $-CH_2OH$, $-CH_2CH_2OH$, $-OH$, $-OCH_3$, $-CH_2OCONH_2$, $-CH_2CH_2COOCH_3$, $-COOCH_3$, $-CH_3$ or $-(CH_2)_3OH$.
- 20 8. Compounds according to any one of claims 1 to 7,
wherein n is 0 or 1.
9. Compounds according to any one of claims 1 to 8,
25 wherein R^5 is an aralkyl group or a heteroaralkyl group.
10. Compounds according to any one of claims 1 to 9,
30 wherein R^5 is a group of formula $-Y-Cy$, Y being a C_1-C_6 alkylene, C_2-C_6 alkenylene or C_1-C_6 heteroalkylene group, wherein optionally a hydrogen atom may have been replaced by a hydroxy group or two hydrogen

atoms may have been replaced by an =O group, and Cy being an optionally substituted phenyl, naphthyl or heteroaryl group containing 1 or 2 rings and from 5 to 10 ring atoms, or an optionally substituted
5 arylheterocycloalkyl or heteroarylheterocycloalkyl group containing two rings and 9 or 10 ring atoms.


11. Compounds according to any one of claims 1 to 10, wherein R^3 is selected from the following groups:



12. Pharmaceutical compositions that comprise a compound according to any one of claims 1 to 11 and, optionally, carrier substances and/or adjuvants.

15

13. Use of a compound or of a pharmaceutical composition according to any one of claims 1 to 12 in the treatment of bacterial infections.



(I)